Full wwPDB X-ray Structure Validation Report (i)

Apr 27, 2024 – 05:01 pm BST

PDB ID	:	5A8J
Title	:	Crystal structure of the ArnB paralog VWA2 from Sulfolobus acidocaldarius
Authors	:	Hoffmann, L.; Anders, K.; Reimann, J.; Linne, U.; Essen, L.-O.; Albers, S.-V.
Deposited on	:	2015-07-16
Resolution	:	1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 6460 atoms, of which 3115 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VWA2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	365	6049	1890	3083	479	590	7	35	26	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q4J9H5
A	2	ALA	-	expression tag	UNP Q4J9H5
A	362	GLY	-	expression tag	UNP Q4J9H5
A	363	SER	-	expression tag	UNP Q4J9H5
A	364	ILE	-	expression tag	UNP Q4J9H5
A	365	GLU	-	expression tag	UNP Q4J9H5
A	366	GLY	-	expression tag	UNP Q4J9H5
A	367	ARG	-	expression tag	UNP Q4J9H5
A	368	HIS	-	expression tag	UNP Q4J9H5
A	369	HIS	-	expression tag	UNP Q4J9H5
A	370	HIS	-	expression tag	UNP Q4J9H5
A	371	HIS	-	expression tag	UNP Q4J9H5
A	372	HIS	-	expression tag	UNP Q4J9H5
A	373	HIS	-	expression tag	UNP Q4J9H5

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 14 3 8 3	0	0
3	A	1	Total C H O 14 3 8 3	0	0
3	A	1	Total C H O 14 3 8 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	14	3	8	3	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
4	A	2	2	2	0	0

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	13	6	7	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
6	A	1	1	1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	323	334	334	0	11

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: VWA2

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	69.64Å 74.28Å 145.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.66 – 1.46 41.66 – 1.46	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.66-1.46) 99.1 (41.66-1.46)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.50 (at 1.46Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.165 , 0.201 0.166 , 0.202	Depositor DCC
R_{free} test set	1340 reflections (2.06%)	wwPDB-VP
Wilson B-factor (Å^2)	16.8	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.43 , 51.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	6460	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, CL, CIT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.52	0/3089	0.69	1/4187 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	142	ASP	CB-CG-OD1	5.75	123.47	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2966	3083	3097	22	1
2	A	5	0	0	0	0
3	A	24	32	32	1	0
4	A	2	0	0	0	0
5	A	13	0	5	2	0
6	A	1	0	0	0	0
7	A	334	0	0	7	0
All	All	3345	3115	3134	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85[A]:ASN:OD1	7:A:2104:HOH:O	1.98	0.82
5:A:1373:CIT:H42	5:A:1373:CIT:O2	1.86	0.74
1:A:227:PRO:HA	7:A:2219:HOH:O	2.03	0.59
1:A:230:ILE:HG21	7:A:2035:HOH:O	2.02	0.58
1:A:225:GLU:OE2	3:A:1369:GOL:O1	2.23	0.57
5:A:1373:CIT:O2	5:A:1373:CIT:C4	2.54	0.56
1:A:315:SER:O	1:A:319[B]:THR:HG23	2.06	0.56
1:A:147:ARG:HA	1:A:150:GLU:HG3	1.91	0.52
1:A:316:GLU:O	1:A:319[A]:THR:HG22	2.10	0.52
1:A:54:LYS:NZ	7:A:2159:HOH:O	2.43	0.51
1:A:85[A]:ASN:ND2	7:A:2104:HOH:O	2.46	0.49
1:A:9[A]:LYS:HE3	7:A:2117:HOH:O	2.13	0.48
1:A:150:GLU:HG2	1:A:176[B]:ILE:CD1	2.47	0.45
1:A:31:PRO:HB3	1:A:207:ALA:CB	2.47	0.44
1:A:37:LEU:HD13	1:A:206[A]:VAL:HG11	1.99	0.44
1:A:206[B]:VAL:HG22	1:A:234:ASP:HB3	2.00	0.43
1:A:171:GLU:HB3	1:A:349:LYS:HG2	2.01	0.42
1:A:306:TYR:CE1	1:A:322[B]:VAL:HG23	2.55	0.42
1:A:85[A]:ASN:CG	7:A:2104:HOH:O	2.49	0.42
1:A:42:TYR:CD1	1:A:72:ILE:HD13	2.54	0.41
1:A:160:LEU:HB3	1:A:181:GLY:HA3	2.03	0.41
1:A:194:LEU:HB3	1:A:195:PRO:HD3	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:ASN:HD22	1:A:244:ASN:HD22[3_655]	1.33	0.27

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/373 (105%)	385 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/339 (105%)	351 (99%)	5 (1%)	67	37

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25[A]	LEU
1	A	25[B]	LEU
1	A	147	ARG
1	A	185	PHE
1	A	288	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GOL	A	1370	-	5,5,5	0.43	0	5,5,5	0.35	0
2	SO4	A	1367	-	4,4,4	0.19	0	6,6,6	0.38	0
3	GOL	A	1369	-	5,5,5	0.32	0	5,5,5	0.45	0
3	GOL	A	1371	-	5,5,5	0.36	0	5,5,5	0.26	0
3	GOL	A	1368	-	5,5,5	0.39	0	5,5,5	0.48	0
5	CIT	A	1373	-	12,12,12	0.97	0	17,17,17	1.71	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1370	-	-	0/4/4/4	-
3	GOL	A	1369	-	-	2/4/4/4	-
3	GOL	A	1371	-	-	0/4/4/4	-
3	GOL	A	1368	-	-	0/4/4/4	-
5	CIT	A	1373	-	-	12/16/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	1373	CIT	O6-C6-C3	4.41	120.70	113.05
5	A	1373	CIT	O2-C1-C2	2.50	122.39	114.35

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1369	GOL	C1-C2-C3-O3
5	A	1373	CIT	C1-C2-C3-O7
5	A	1373	CIT	C1-C2-C3-C4
5	A	1373	CIT	C1-C2-C3-C6
5	A	1373	CIT	O7-C3-C6-O5
5	A	1373	CIT	O7-C3-C6-O6
5	A	1373	CIT	C4-C3-C6-O5
5	A	1373	CIT	C4-C3-C6-O6
3	A	1369	GOL	O2-C2-C3-O3
5	A	1373	CIT	C2-C3-C6-O5
5	A	1373	CIT	C2-C3-C6-O6
5	A	1373	CIT	C2-C3-C4-C5
5	A	1373	CIT	C6-C3-C4-C5
5	A	1373	CIT	O2-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1369	GOL	1	0
5	A	1373	CIT	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	365/373 (97%)	0.05	9 (2%) 57 60	8, 15, 32, 43	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	VAL	4.1
1	A	333	LYS	3.7
1	A	346	ASN	3.7
1	A	338	ILE	2.8
1	A	366	GLY	2.5
1	A	87	ILE	2.4
1	A	313	GLY	2.3
1	A	332	THR	2.3
1	A	152	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	GOL	A	1369	6/6	0.76	0.25	46,55,57,57	0
3	GOL	A	1371	6/6	0.83	0.21	35,42,48,51	0
5	CIT	A	1373	13/13	0.90	0.12	21,25,26,27	13
3	GOL	A	1370	6/6	0.91	0.24	44,53,55,55	0
4	CL	A	1372	1/1	0.92	0.08	59,59,59,59	0
4	CL	A	1375	1/1	0.93	0.17	50,50,50,50	1
6	NA	A	1374	1/1	0.93	0.21	43,43,43,43	0
3	GOL	A	1368	6/6	0.94	0.12	20,25,29,30	0
2	SO4	A	1367	5/5	0.98	0.08	28,30,32,35	0

6.5 Other polymers (i)

There are no such residues in this entry.