wwPDB X-ray Structure Validation Summary Report (i)

Apr 27, 2024 – 04:05 pm BST

PDB ID	:	4A9A
Title	:	Structure of Rbg1 in complex with Tma46 dfrp domain
Authors	:	Francis, S.M.; Gas, M.; Daugeron, M.; Seraphin, B.; Bravo, J.
Deposited on	:	2011-11-25
Resolution	:	2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	
1	B	376	
2	C	142	
2	D	142	

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBOSOME-INTERACTING GTPASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2776	1764	480	523	9	0	0	0
1	B	355	2738	1736	476	517	9	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP P39729
A	-5	HIS	-	expression tag	UNP P39729
A	-4	HIS	-	expression tag	UNP P39729
A	-3	HIS	-	expression tag	UNP P39729
A	-2	HIS	-	expression tag	UNP P39729
A	-1	HIS	-	expression tag	UNP P39729
A	0	HIS	-	expression tag	UNP P39729
B	-6	MET	-	expression tag	UNP P39729
B	-5	HIS	-	expression tag	UNP P39729
B	-4	HIS	-	expression tag	UNP P39729
B	-3	HIS	-	expression tag	UNP P39729
B	-2	HIS	-	expression tag	UNP P39729
B	-1	HIS	-	expression tag	UNP P39729
B	0	HIS	-	expression tag	UNP P39729

- Molecule 2 is a protein called TRANSLATION MACHINERY-ASSOCIATED PROTEIN 46.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	106	844	536	145	162	1	0	0	0
2	D	86	672	427	117	127	1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	204	MET	-	expression tag	UNP Q12000
D	204	MET	-	expression tag	UNP Q12000

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	36	36	36	0	0
			Total	O		
3	B	24	24	24	0	0
			Total	O		
3	C	13	13	13	0	0
			Total	O		
3	D	4	4	4	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RIBOSOME-INTERACTING GTPASE 1

Chain A:



• Molecule 1: RIBOSOME-INTERACTING GTPASE 1

Chain B:



• Molecule 2: TRANSLATION MACHINERY-ASSOCIATED PROTEIN 46

Chain C:



• Molecule 2: TRANSLATION MACHINERY-ASSOCIATED PROTEIN 46

Chain D:

5%
52%
8%
39%


MET GLU GLN LYS ARG LEU GLU ARG GLU SER L214 E215 K216 F225 I226 E227 T228 E229 I258 K262 K263 L264 S265 S266 K267 R268 I276 M279 S280 A281 E282 ASN LYS SER PHE GLU THR ASP ASN ALA ASP MET PRO ASP ASP VAL THR GLN GLY SER ALA TRP ASP LEU THR

GLU PHE THR ASP ALA LEU LYS LYS ALA ASP HIS GLN ASP D320 G321 I336 LYS LYS ALA ASN SER ALA THR LEU ALA

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	86.20Å 224.89Å 84.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.22 – 2.67 56.22 – 2.67	Depositor EDS
% Data completeness (in resolution range)	100.0 (56.22-2.67) 100.0 (56.22-2.67)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.68 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.196 , 0.222 0.196 , 0.222	Depositor DCC
R_{free} test set	1200 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å^2)	65.8	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 43.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.025 for l,-k,h	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	7107	wwPDB-VP
Average B, all atoms (Å^2)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.47	0/2817	0.62	0/3807
1	B	0.45	0/2777	0.60	0/3751
2	C	0.51	1/857 (0.1%)	0.62	0/1146
2	D	0.47	0/681	0.60	0/909
All	All	0.47	1/7132 (0.0%)	0.61	0/9613

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	303	TRP	CD2-CE2	5.03	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2776	0	2869	32	0
1	B	2738	0	2826	23	0
2	C	844	0	865	6	0
2	D	672	0	695	4	0
3	A	36	0	0	0	0
3	B	24	0	0	1	0
3	C	13	0	0	0	0
3	D	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7107	0	7255	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:258:ILE:O	2:D:262:LYS:HG2	1.80	0.82
1:B:58:PHE:HB2	1:B:142:THR:HG21	1.64	0.80
1:B:58:PHE:HB2	1:B:142:THR:CG2	2.11	0.80
1:A:68:VAL:HG11	1:A:147:PHE:HE2	1.54	0.73
1:B:147:PHE:HE1	1:B:246:ILE:HD12	1.55	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/376 (93%)	337 (96%)	12 (3%)	1 (0%)	41 64
1	B	349/376 (93%)	331 (95%)	14 (4%)	4 (1%)	14 31
2	C	102/142 (72%)	97 (95%)	5 (5%)	0	100 100
2	D	82/142 (58%)	76 (93%)	3 (4%)	3 (4%)	3 6
All	All	883/1036 (85%)	841 (95%)	34 (4%)	8 (1%)	17 37

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	46	ALA
1	B	120	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	204	ASP
2	D	265	SER
1	A	307	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/325 (95%)	297 (96%)	13 (4%)	30	55
1	B	305/325 (94%)	294 (96%)	11 (4%)	35	61
2	C	91/123 (74%)	90 (99%)	1 (1%)	73	89
2	D	72/123 (58%)	69 (96%)	3 (4%)	30	55
All	All	778/896 (87%)	750 (96%)	28 (4%)	35	61

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	142	THR
2	D	276	ILE
1	B	214	ARG
2	C	305	LEU
1	B	207	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	325	ASN
1	A	350	GLN
1	B	325	ASN
1	B	194	ASN
1	A	216	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	356/376 (94%)	0.24	5 (1%)	75	76	35, 65, 121, 173	0
1	B	355/376 (94%)	0.05	8 (2%)	60	60	42, 71, 127, 186	0
2	C	106/142 (74%)	0.12	4 (3%)	40	38	39, 75, 132, 193	0
2	D	86/142 (60%)	0.50	7 (8%)	12	10	53, 94, 169, 193	0
All	All	903/1036 (87%)	0.18	24 (2%)	54	54	35, 71, 132, 193	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	214	LEU	6.0
2	D	216	LYS	5.8
2	C	215	GLU	4.7
1	B	197	PRO	4.0
1	B	52	GLY	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.