



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 27, 2024 – 11:50 pm BST

PDB ID : 1HF2
Title : Crystal structure of the bacterial cell-division inhibitor MinC from *T. maritima*
Authors : Cordell, S.C.; Anderson, R.E.; Lowe, J.
Deposited on : 2000-11-27
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<div> <div>8%</div> <div>54%</div> <div>36%</div> <div>7%</div> </div>
1	B	210	<div> <div>%</div> <div>60%</div> <div>35%</div> <div>..</div> </div>
1	C	210	<div> <div>6%</div> <div>61%</div> <div>32%</div> <div>..</div> </div>
1	D	210	<div> <div>17%</div> <div>51%</div> <div>35%</div> <div>10%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6685 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SEPTUM SITE-DETERMINING PROTEIN MINC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1487	936	270	277	4			
1	B	206	Total	C	N	O	S	0	0	0
			1563	980	282	298	3			
1	C	202	Total	C	N	O	S	0	0	0
			1532	963	276	289	4			
1	D	190	Total	C	N	O	S	0	0	0
			1443	908	262	270	3			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	139	Total	O	0	0
			139	139		
2	B	202	Total	O	0	0
			202	202		
2	C	174	Total	O	0	0
			174	174		
2	D	145	Total	O	0	0
			145	145		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SEPTUM SITE-DETERMINING PROTEIN MINC



• Molecule 1: SEPTUM SITE-DETERMINING PROTEIN MINC



• Molecule 1: SEPTUM SITE-DETERMINING PROTEIN MINC



K203	V204	S205	F206	GLU	ARG	SER	GLU
------	------	------	------	-----	-----	-----	-----

● Molecule 1: SEPTUM SITE-DETERMINING PROTEIN MINC



MET	VAL	D3	F4	K5	E9	G10	L11	V12	L13	L14	I15	K16	D17	Y18	Q19	N20	L21	E22	E23	V24	L25	N26	A27	I28	S29	A30	R31	I32	T33	Q34	M35	G36	G37	F38	F39	A40	K41	G42	D43	R44	I45	E50	N51	H52	N53	K54	H55	P60	R61	I62	V63	S64	H65	L66	R67	N68
-----	-----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

L69	G70	L71	E72	V73	S74	Q75	I76	L77	V78	GLY	SER	THR	VAL	GLU	GLY	LYS	GLU	ASN	ASP	LEU	LYS	VAL	GLN	S93	R94	T95	T96	V97	E98	S99	T100	G101	K102	R106	R109	S110	G111	Q112	S117	G118	D119	N127	L133	A134	G135	F141	Q145	G146	N147	I148	R149
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

V159	V160	Q166	T167	S168	L169	T170	Q171	I176	T177	E182	P186	S187	I188	G193	N194	R195	F201	V204	S205	F206	GLU	ARG	SER	GLU
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.58Å 106.09Å 162.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.20 36.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.8 (100.00-2.20) 93.5 (36.98-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.98 (at 2.10Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.237 , 0.300 0.241 , 0.233	Depositor DCC
R_{free} test set	2515 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.974	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 62.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6685	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1503	0.61	0/2021
1	B	0.34	0/1580	0.64	0/2127
1	C	0.35	0/1548	0.66	0/2083
1	D	0.33	0/1459	0.63	0/1963
All	All	0.34	0/6090	0.64	0/8194

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1487	0	1543	71	0
1	B	1563	0	1608	65	0
1	C	1532	0	1585	74	0
1	D	1443	0	1489	86	0
2	A	139	0	0	15	0
2	B	202	0	0	13	0
2	C	174	0	0	19	0
2	D	145	0	0	19	0
All	All	6685	0	6225	294	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 294 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:59:ILE:HG13	1:C:60:PRO:HD3	1.43	1.00
1:D:188:ILE:HD12	1:D:204:VAL:HG21	1.43	1.00
1:C:75:GLN:HE22	1:C:94:ARG:HE	1.11	0.94
1:B:68:ASN:HB3	1:C:61:ARG:HD3	1.49	0.93
1:C:149:ARG:HH11	1:C:171:GLN:NE2	1.67	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/210 (91%)	174 (91%)	12 (6%)	6 (3%)	4	2
1	B	204/210 (97%)	189 (93%)	14 (7%)	1 (0%)	29	31
1	C	198/210 (94%)	178 (90%)	15 (8%)	5 (2%)	5	3
1	D	186/210 (89%)	165 (89%)	16 (9%)	5 (3%)	5	2
All	All	780/840 (93%)	706 (90%)	57 (7%)	17 (2%)	6	4

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	50	GLU
1	B	194	ASN
1	C	50	GLU
1	D	43	ASP
1	D	100	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/176 (93%)	153 (94%)	10 (6%)	18	21
1	B	172/176 (98%)	164 (95%)	8 (5%)	26	33
1	C	169/176 (96%)	161 (95%)	8 (5%)	26	33
1	D	158/176 (90%)	148 (94%)	10 (6%)	18	20
All	All	662/704 (94%)	626 (95%)	36 (5%)	22	26

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	23	GLU
1	D	195	ARG
1	D	34	GLN
1	D	51	ASN
1	B	51	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	57	GLN
1	D	20	ASN
1	C	65	HIS
1	C	127	ASN
1	D	51	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/210 (93%)	0.51	16 (8%) 11 10	24, 45, 66, 80	0
1	B	206/210 (98%)	0.16	3 (1%) 73 72	18, 34, 59, 73	0
1	C	202/210 (96%)	0.51	13 (6%) 19 18	17, 39, 68, 84	0
1	D	190/210 (90%)	0.98	36 (18%) 1 1	15, 43, 94, 105	0
All	All	794/840 (94%)	0.53	68 (8%) 10 9	15, 40, 73, 105	0

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	38	PHE	7.3
1	D	33	THR	6.8
1	D	21	LEU	6.2
1	D	30	ALA	5.7
1	D	28	ILE	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.