



Full wwPDB X-ray Structure Validation Report i

Sep 20, 2023 – 09:33 PM EDT

PDB ID : 5IQE
Title : Aminoglycoside Phosphotransferase (2")-Ia (CTD of AAC(6')-Ie/APH(2")-Ia) in complex with GMPPNP, Magnesium, and Neomycin B
Authors : Caldwell, S.J.; Berghuis, A.M.
Deposited on : 2016-03-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 10539 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional AAC/APH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	2423	1547	376	490	10	0	0	0
1	B	295	2425	1551	374	490	10	0	0	0
1	C	292	2410	1540	373	487	10	0	0	0
1	D	292	2411	1540	372	489	10	0	0	0

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	32	10	6	13	3	0	0
2	B	1	41	10	7	19	5	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C N O P 32 10 6 13 3	0	0
2	D	1	Total C N O P 32 10 6 13 3	0	0

- Molecule 3 is NEOMYCIN (three-letter code: NMY) (formula: C₂₃H₄₆N₆O₁₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 42 23 6 13	0	0
3	B	1	Total C N O 42 23 6 13	0	0
3	C	1	Total C N O 42 23 6 13	0	0
3	D	1	Total C N O 42 23 6 13	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Mg 2 2	0	0
4	B	2	Total Mg 2 2	0	0
4	C	2	Total Mg 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	2	Total Mg 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	147	Total O 147 147	0	0
6	B	167	Total O 167 167	0	0
6	C	138	Total O 138 138	0	0
6	D	99	Total O 99 99	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional AAC/APH




- Molecule 1: Bifunctional AAC/APH



- Molecule 1: Bifunctional AAC/APH



- Molecule 1: Bifunctional AAC/APH



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.18Å 100.36Å 93.94Å 90.00° 105.30° 90.00°	Depositor
Resolution (Å)	90.61 – 2.50 65.73 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (90.61-2.50) 100.0 (65.73-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.04 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R , R_{free}	0.169 , 0.214 0.176 , 0.217	Depositor DCC
R_{free} test set	2804 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	54.6	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.9	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10539	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GNP, NMY, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2466	0.75	0/3327
1	B	0.69	1/2469 (0.0%)	0.74	2/3331 (0.1%)
1	C	0.70	1/2453 (0.0%)	0.75	0/3306
1	D	0.59	0/2453	0.71	0/3308
All	All	0.66	2/9841 (0.0%)	0.74	2/13272 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	377	CYS	CB-SG	-5.08	1.73	1.81
1	C	466	GLU	CG-CD	5.07	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	377	CYS	CB-CA-C	-5.75	98.91	110.40
1	B	301	ASP	CB-CG-OD2	-5.08	113.73	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2423	0	2327	15	0
1	B	2425	0	2323	14	0
1	C	2410	0	2322	13	0
1	D	2411	0	2313	9	0
2	A	32	0	13	1	0
2	B	41	0	2	1	0
2	C	32	0	13	1	0
2	D	32	0	13	1	0
3	A	42	0	46	7	0
3	B	42	0	46	6	0
3	C	42	0	46	2	0
3	D	42	0	46	3	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	B	6	0	8	0	0
6	A	147	0	0	7	0
6	B	167	0	0	4	0
6	C	138	0	0	8	0
6	D	99	0	0	4	0
All	All	10539	0	9518	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:500[B]:GNP:O1B	6:B:904:HOH:O	1.72	1.05
1:A:230:SER:HB2	1:A:267:ASP:O	1.54	1.05
1:D:230:SER:HB2	1:D:267:ASP:O	1.65	0.94
1:B:230:SER:HB2	1:B:267:ASP:O	1.66	0.93
1:A:319:GLU:OE2	6:A:1274:HOH:O	2.01	0.78
1:B:374:ASP:OD1	3:B:600:NMY:N7	2.17	0.78
1:D:254:ASN:HB2	6:D:1500:HOH:O	1.87	0.73
2:A:500:GNP:O1B	6:A:901:HOH:O	2.08	0.72
6:A:1274:HOH:O	1:C:265:ILE:O	2.10	0.68
1:A:230:SER:CB	1:A:267:ASP:O	2.35	0.68
1:A:267:ASP:HB2	6:D:1123:HOH:O	1.96	0.65
1:D:307:ARG:NH2	6:D:1005:HOH:O	2.30	0.64
1:C:185:ASN:ND2	1:D:184:THR:OG1	2.31	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:296:ASN:ND2	6:D:1097:HOH:O	2.31	0.62
1:C:298:LEU:HA	6:C:1268:HOH:O	2.03	0.58
1:A:451:GLU:OE2	3:A:600:NMY:N23	2.38	0.57
1:B:267:ASP:HB2	6:B:1058:HOH:O	2.04	0.57
3:A:600:NMY:N2	6:A:1151:HOH:O	2.31	0.57
1:B:298:LEU:HA	6:B:1268:HOH:O	2.05	0.56
1:A:319:GLU:HB3	6:A:1317:HOH:O	2.08	0.54
1:C:228:LYS:HE3	6:C:1038:HOH:O	2.07	0.54
1:A:230:SER:HB3	1:A:268:GLU:HA	1.91	0.53
3:A:600:NMY:H1	3:A:600:NMY:C13	2.39	0.53
1:B:211:GLY:HA3	6:B:1091:HOH:O	2.09	0.53
6:C:1220:HOH:O	1:D:184:THR:HG22	2.09	0.52
3:D:600:NMY:H1	3:D:600:NMY:H13	1.90	0.52
1:B:189:MET:SD	1:B:227:THR:HG21	2.50	0.52
3:B:600:NMY:H13	3:B:600:NMY:H1	1.92	0.52
1:C:413:SER:HB2	3:C:600:NMY:N6	2.25	0.52
1:A:185:ASN:OD1	1:B:179:TYR:OH	2.26	0.51
3:C:600:NMY:H13	3:C:600:NMY:H2	1.92	0.51
1:C:427:ARG:NH2	6:C:1292:HOH:O	2.43	0.51
1:B:189:MET:HG2	1:B:217:TYR:CE1	2.46	0.51
1:B:451:GLU:OE2	3:B:600:NMY:N23	2.43	0.50
1:C:189:MET:HG2	1:C:217:TYR:CE1	2.47	0.50
1:A:374:ASP:OD1	3:A:600:NMY:N7	2.44	0.50
1:A:189:MET:SD	1:A:227:THR:HG21	2.52	0.50
3:A:600:NMY:H1	3:A:600:NMY:H13	1.94	0.49
1:B:286:ILE:O	1:B:289:THR:HB	2.13	0.49
1:C:325:LYS:NZ	6:C:1258:HOH:O	2.31	0.48
1:C:395:GLY:HA3	6:C:902:HOH:O	2.13	0.48
1:D:189:MET:SD	1:D:227:THR:HG21	2.53	0.48
2:C:500:GNP:O2B	6:C:1221:HOH:O	2.20	0.47
1:C:189:MET:SD	1:C:227:THR:HG21	2.55	0.47
3:B:600:NMY:H231	3:B:600:NMY:O18	2.15	0.46
1:A:324:ASN:OD1	6:A:980:HOH:O	2.20	0.46
1:A:415:GLU:OE2	3:A:600:NMY:H62	2.16	0.46
1:C:228:LYS:CE	6:C:1038:HOH:O	2.63	0.46
1:D:186:VAL:HG13	1:D:205:ILE:HG23	1.98	0.46
1:B:198:ASP:OD1	1:B:199:ASN:N	2.42	0.45
3:B:600:NMY:H1	3:B:600:NMY:C13	2.46	0.45
1:A:189:MET:HG2	1:A:217:TYR:CE1	2.52	0.45
1:B:415:GLU:OE2	3:B:600:NMY:H62	2.17	0.45
1:A:186:VAL:HG13	1:A:205:ILE:HG23	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:189:MET:HG2	1:D:217:TYR:CE2	2.52	0.44
3:D:600:NMY:H1	3:D:600:NMY:C13	2.47	0.44
3:A:600:NMY:H232	3:A:600:NMY:O18	2.17	0.44
2:D:500:GNP:O1G	3:D:600:NMY:N19	2.47	0.44
1:A:319:GLU:CB	6:A:1317:HOH:O	2.65	0.43
1:B:186:VAL:HG13	1:B:205:ILE:HG23	2.00	0.43
1:C:186:VAL:HG13	1:C:205:ILE:HG23	2.00	0.43
1:C:252:GLU:O	1:C:308:GLN:NE2	2.46	0.43
1:B:237:TYR:HB2	1:B:270:SER:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	291/305 (95%)	278 (96%)	11 (4%)	2 (1%)	22 39
1	B	291/305 (95%)	279 (96%)	11 (4%)	1 (0%)	41 61
1	C	288/305 (94%)	276 (96%)	11 (4%)	1 (0%)	41 61
1	D	286/305 (94%)	276 (96%)	10 (4%)	0	100 100
All	All	1156/1220 (95%)	1109 (96%)	43 (4%)	4 (0%)	41 61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	230	SER
1	A	198	ASP
1	B	198	ASP
1	C	198	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	268/281 (95%)	261 (97%)	7 (3%)	46 72
1	B	267/281 (95%)	261 (98%)	6 (2%)	52 77
1	C	267/281 (95%)	261 (98%)	6 (2%)	52 77
1	D	268/281 (95%)	263 (98%)	5 (2%)	57 80
All	All	1070/1124 (95%)	1046 (98%)	24 (2%)	52 77

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198	ASP
1	A	221	ASN
1	A	230	SER
1	A	265	ILE
1	A	267	ASP
1	A	308	GLN
1	A	317	ILE
1	B	221	ASN
1	B	230	SER
1	B	265	ILE
1	B	267	ASP
1	B	285	GLU
1	B	289	THR
1	C	198	ASP
1	C	221	ASN
1	C	265	ILE
1	C	267	ASP
1	C	387	ARG
1	C	479	ASP
1	D	221	ASN
1	D	230	SER
1	D	265	ILE
1	D	267	ASP
1	D	387	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	341	ASN
1	C	327	ASN
1	C	341	ASN
1	D	295	GLN
1	D	296	ASN
1	D	420	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 18 ligands modelled in this entry, 8 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNP	D	500	4	29,34,34	1.38	3 (10%)	33,54,54	2.27	8 (24%)
2	GNP	A	500	4	29,34,34	1.77	5 (17%)	33,54,54	2.14	9 (27%)
3	NMY	D	600	-	45,45,45	0.35	0	63,67,67	1.32	8 (12%)
3	NMY	C	600	-	45,45,45	0.31	0	63,67,67	1.48	6 (9%)
3	NMY	A	600	-	45,45,45	0.27	0	63,67,67	1.51	4 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	804	-	5,5,5	0.85	0	5,5,5	0.95	0
2	GNP	B	500[B]	4	29,34,34	0.95	2 (6%)	33,54,54	2.08	6 (18%)
2	GNP	B	500[A]	4	29,34,34	0.97	2 (6%)	33,54,54	2.21	9 (27%)
3	NMY	B	600	-	45,45,45	0.29	0	63,67,67	1.51	6 (9%)
2	GNP	C	500	4	29,34,34	0.91	2 (6%)	33,54,54	2.18	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNP	D	500	4	-	1/14/38/38	0/3/3/3
2	GNP	A	500	4	-	1/14/38/38	0/3/3/3
3	NMY	D	600	-	-	9/18/94/94	0/4/4/4
3	NMY	C	600	-	-	4/18/94/94	0/4/4/4
3	NMY	A	600	-	-	9/18/94/94	0/4/4/4
5	GOL	B	804	-	-	1/4/4/4	-
2	GNP	B	500[B]	4	-	1/14/38/38	0/3/3/3
2	GNP	B	500[A]	4	-	2/14/38/38	0/3/3/3
3	NMY	B	600	-	-	7/18/94/94	0/4/4/4
2	GNP	C	500	4	-	4/14/38/38	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	GNP	PG-O2G	4.48	1.68	1.56
2	A	500	GNP	PG-O3G	4.47	1.68	1.56
2	D	500	GNP	PG-O3G	4.27	1.68	1.56
2	A	500	GNP	PB-O3A	-4.17	1.53	1.59
2	D	500	GNP	C6-N1	3.44	1.39	1.33
2	A	500	GNP	C6-N1	3.42	1.39	1.33
2	C	500	GNP	C6-N1	2.75	1.37	1.33
2	B	500[A]	GNP	C6-N1	2.73	1.37	1.33
2	B	500[B]	GNP	C6-N1	2.73	1.37	1.33
2	D	500	GNP	PB-O2B	2.66	1.63	1.56
2	B	500[B]	GNP	PG-O2G	2.43	1.63	1.56
2	B	500[A]	GNP	PB-O2B	2.27	1.62	1.56
2	A	500	GNP	PA-O5'	2.15	1.68	1.59
2	C	500	GNP	PG-O2G	2.07	1.62	1.56

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	500	GNP	C5-C6-N1	-8.25	112.15	123.43
3	B	600	NMY	O12-C12-C7	7.90	124.28	109.81
2	B	500[A]	GNP	C5-C6-N1	-7.83	112.72	123.43
2	B	500[B]	GNP	C5-C6-N1	-7.83	112.72	123.43
3	A	600	NMY	O12-C12-C7	7.80	124.11	109.81
2	A	500	GNP	C5-C6-N1	-7.80	112.76	123.43
2	D	500	GNP	C5-C6-N1	-7.58	113.07	123.43
3	C	600	NMY	O11-C13-O16	-6.74	104.13	111.43
3	A	600	NMY	O11-C13-O16	-6.20	104.71	111.43
2	C	500	GNP	C2-N1-C6	6.04	125.52	115.93
2	D	500	GNP	O2B-PB-O1B	-5.91	97.53	109.92
2	B	500[A]	GNP	C2-N1-C6	5.74	125.05	115.93
2	B	500[B]	GNP	C2-N1-C6	5.74	125.05	115.93
2	D	500	GNP	C2-N1-C6	5.54	124.73	115.93
2	A	500	GNP	C2-N1-C6	5.20	124.20	115.93
3	B	600	NMY	O11-C13-O16	-5.10	105.91	111.43
3	C	600	NMY	O12-C12-C7	4.29	117.67	109.81
3	C	600	NMY	C11-C12-C7	3.92	119.20	109.63
3	D	600	NMY	O12-C12-C11	3.78	119.97	109.94
3	D	600	NMY	O12-C12-C7	3.57	116.36	109.81
3	D	600	NMY	C11-C12-C7	3.42	117.98	109.63
3	C	600	NMY	O12-C12-C11	3.24	118.53	109.94
3	D	600	NMY	O11-C13-O16	-3.20	107.96	111.43
2	D	500	GNP	N3-C2-N1	-3.16	123.01	127.22
2	A	500	GNP	N3-C2-N1	-3.11	123.07	127.22
2	D	500	GNP	O2G-PG-O1G	-3.04	105.82	113.45
3	B	600	NMY	O1-C1-C2	3.03	113.44	108.22
2	B	500[A]	GNP	O2G-PG-O1G	-3.00	105.90	113.45
2	B	500[B]	GNP	O2B-PB-O1B	-2.98	103.67	109.92
2	C	500	GNP	N3-C2-N1	-2.96	123.28	127.22
3	C	600	NMY	O5-C1-C2	2.91	116.61	110.06
2	B	500[A]	GNP	N3-C2-N1	-2.81	123.47	127.22
2	B	500[B]	GNP	N3-C2-N1	-2.81	123.47	127.22
2	A	500	GNP	C1'-N9-C4	-2.81	121.70	126.64
3	D	600	NMY	O1-C1-C2	2.73	112.92	108.22
2	D	500	GNP	C2-N3-C4	-2.72	112.25	115.36
3	D	600	NMY	O18-C18-O22	-2.64	103.30	110.67
2	B	500[A]	GNP	C2-N3-C4	-2.63	112.35	115.36
2	B	500[B]	GNP	C2-N3-C4	-2.63	112.35	115.36
2	A	500	GNP	O2B-PB-O1B	-2.62	104.42	109.92
2	C	500	GNP	C2-N3-C4	-2.53	112.47	115.36
3	B	600	NMY	C9-C8-C7	-2.49	106.06	111.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	500	GNP	C4-C5-C6	-2.48	118.43	120.80
2	A	500	GNP	O2'-C2'-C1'	-2.48	101.70	110.85
3	A	600	NMY	C9-C8-C7	-2.44	106.17	111.18
2	B	500[A]	GNP	O2B-PB-O1B	-2.44	104.81	109.92
2	C	500	GNP	O1G-PG-N3B	2.41	115.32	111.77
2	B	500[A]	GNP	C4-C5-C6	-2.41	118.50	120.80
2	B	500[B]	GNP	C4-C5-C6	-2.41	118.50	120.80
2	D	500	GNP	O1G-PG-N3B	-2.40	108.23	111.77
2	A	500	GNP	O3'-C3'-C2'	-2.38	104.11	111.82
2	C	500	GNP	O2B-PB-O1B	-2.37	104.94	109.92
2	A	500	GNP	C2-N3-C4	-2.30	112.73	115.36
2	B	500[A]	GNP	O3G-PG-O1G	-2.23	107.83	113.45
2	D	500	GNP	C4-C5-C6	-2.22	118.68	120.80
3	B	600	NMY	O1-C10-C9	-2.22	103.88	109.18
3	B	600	NMY	O1-C10-C11	2.22	113.12	107.48
3	D	600	NMY	O22-C22-C21	2.19	113.67	109.69
3	A	600	NMY	O22-C22-C23	2.16	110.03	106.01
2	B	500[A]	GNP	O1G-PG-N3B	2.15	114.93	111.77
3	D	600	NMY	O1-C10-C9	-2.13	104.11	109.18
3	C	600	NMY	C9-C8-C7	-2.12	106.84	111.18
2	A	500	GNP	C4-C5-C6	-2.01	118.88	120.80

There are no chirality outliers.

All (39) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	500[A]	GNP	PB-N3B-PG-O1G
2	B	500[A]	GNP	PG-N3B-PB-O1B
2	C	500	GNP	PB-N3B-PG-O1G
2	C	500	GNP	PG-N3B-PB-O1B
2	D	500	GNP	PG-N3B-PB-O1B
3	A	600	NMY	O5-C5-C6-N6
3	A	600	NMY	C14-C13-O11-C11
3	A	600	NMY	O16-C13-O11-C11
3	B	600	NMY	C14-C13-O11-C11
3	B	600	NMY	O16-C13-O11-C11
3	B	600	NMY	O22-C22-C23-N19
3	C	600	NMY	C14-C13-O11-C11
3	C	600	NMY	O16-C13-O11-C11
3	C	600	NMY	C14-C15-O18-C18
3	D	600	NMY	C14-C13-O11-C11
3	D	600	NMY	O16-C13-O11-C11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	D	600	NMY	C21-C22-C23-N19
3	D	600	NMY	O22-C22-C23-N19
3	D	600	NMY	O16-C16-C17-O17
3	D	600	NMY	C15-C16-C17-O17
3	B	600	NMY	O16-C16-C17-O17
3	B	600	NMY	C15-C16-C17-O17
3	C	600	NMY	O22-C18-O18-C15
3	A	600	NMY	O16-C16-C17-O17
3	A	600	NMY	C15-C16-C17-O17
2	A	500	GNP	PB-O3A-PA-O1A
3	D	600	NMY	O5-C1-O1-C10
3	B	600	NMY	C11-C10-O1-C1
3	A	600	NMY	C4-C5-C6-N6
3	A	600	NMY	C21-C22-C23-N19
3	D	600	NMY	C11-C10-O1-C1
3	B	600	NMY	O5-C1-O1-C10
5	B	804	GOL	O1-C1-C2-O2
2	C	500	GNP	PG-N3B-PB-O3A
3	A	600	NMY	O5-C1-O1-C10
3	A	600	NMY	C11-C10-O1-C1
2	B	500[B]	GNP	PB-O3A-PA-O2A
2	C	500	GNP	PB-O3A-PA-O2A
3	D	600	NMY	O22-C18-O18-C15


There are no ring outliers.


8 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	500	GNP	1	0
2	A	500	GNP	1	0
3	D	600	NMY	3	0
3	C	600	NMY	2	0
3	A	600	NMY	7	0
2	B	500[B]	GNP	1	0
3	B	600	NMY	6	0
2	C	500	GNP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/305 (96%)	0.17	2 (0%) 87 89	40, 64, 97, 133	0
1	B	295/305 (96%)	0.09	1 (0%) 94 94	40, 62, 97, 126	0
1	C	292/305 (95%)	0.11	1 (0%) 94 94	36, 62, 100, 166	0
1	D	292/305 (95%)	0.25	8 (2%) 54 58	54, 77, 114, 147	0
All	All	1174/1220 (96%)	0.15	12 (1%) 82 84	36, 67, 104, 166	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	THR	4.5
1	D	179	TYR	4.2
1	D	473	TYR	3.3
1	B	179	TYR	3.3
1	D	477	TYR	3.2
1	D	180	ASP	2.4
1	D	254	ASN	2.2
1	C	479	ASP	2.1
1	D	464	PHE	2.1
1	D	181	ASP	2.1
1	A	213	ASP	2.1
1	D	251	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NMY	A	600	42/42	0.62	0.30	153,177,217,242	0
3	NMY	C	600	42/42	0.68	0.27	193,217,282,324	0
3	NMY	B	600	42/42	0.86	0.23	114,121,154,169	0
5	GOL	B	804	6/6	0.89	0.16	72,87,96,99	0
4	MG	A	702	1/1	0.95	0.07	52,52,52,52	0
4	MG	B	702	1/1	0.95	0.08	59,59,59,59	0
4	MG	D	702	1/1	0.95	0.06	59,59,59,59	0
3	NMY	D	600	42/42	0.95	0.16	68,78,89,93	0
2	GNP	B	500[B]	32/32	0.97	0.13	45,47,50,50	9
4	MG	C	700	1/1	0.97	0.10	61,61,61,61	0
4	MG	A	700	1/1	0.97	0.10	51,51,51,51	0
2	GNP	B	500[A]	32/32	0.97	0.13	45,47,52,53	9
2	GNP	D	500	32/32	0.98	0.13	54,61,67,70	0
4	MG	C	702	1/1	0.98	0.06	58,58,58,58	0
4	MG	D	700	1/1	0.98	0.11	48,48,48,48	0
4	MG	B	700	1/1	0.98	0.08	43,43,43,43	0
2	GNP	A	500	32/32	0.98	0.15	39,46,57,63	0
2	GNP	C	500	32/32	0.99	0.15	50,55,61,63	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.