wwPDB X-ray Structure Validation Summary Report (i)

Apr 27, 2024 – 02:15 pm BST

PDB ID	:	2JJK
Title	:	FRUCTOSE-1,6-BISPHOSPHATASE(D-FRUCTOSE-1,6-BISPHOSPHATE -1- PHOSPHOHYDROLASE) (E.C.3.1.3.11) COMPLEXED WITH A DUAL BINDING AMP SITE INHIBITOR
Authors	:	Ruf, A.; Joseph, C.; Benz, J.; Fol, B.; Tetaz, T.; Hebeisen, P.
Deposited on	:	2008-04-09
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	.% 80% 14% • 5%
1	B	338	3% 81% 12% • 6%
1	C	338	.% 80% 14% 6%
1	D	338	.% 78% 16% 6%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 10895 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOSE-1,6-BISPHOSPHATASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	320	2442	1556	410	459	17	0	0	1
1	B	318	2426	1545	407	457	17	0	0	1
1	C	319	2435	1551	409	458	17	0	0	1
1	D	318	2426	1545	407	457	17	0	0	1

- Molecule 2 is N,N'-(heptane-1,7-diyldicarbamoyl)bis(3-chlorobenzenesulfonamide) (three-letter code: R15) (formula: $C_{21}H_{26}Cl_2N_4O_6S_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	N	O	S		
2	A	1	42	28	2	4	6	2	0	1
2	D	1	42	28	2	4	6	2	0	1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	217	217	217	0	0
			Total	O		
3	B	270	270	270	0	0
			Total	O		
3	C	292	292	292	0	0
			Total	O		
3	D	303	303	303	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FRUCTOSE-1,6-BISPHOSPHATASE 1

Chain A:



• Molecule 1: FRUCTOSE-1,6-BISPHOSPHATASE 1

Chain B:



• Molecule 1: FRUCTOSE-1,6-BISPHOSPHATASE 1

Chain C:



• Molecule 1: FRUCTOSE-1,6-BISPHOSPHATASE 1

Chain D:


MET
ALA
ASP
GLN
ALA
PRO
PHE
ASP
THR
D9
V17
M18
G21
R22
R25
G26
T31
Q32
C38
I44
S45
V48
R49
K50
A51
G61
SER
THR
ASN
VAL
THR
GLY
ASP
GLN
VAL
LYS
K72
L73
A90
K100
H101
A102
G111
K112
D121
N125
I126
R140
K141
K142
S143
E146
E149
D178
N182
C183
F184
G191
E192
F193
K198
D199
K204
K205
G206
A216
F219
E225
Y226
I227
K231
N236
G260
Y264
P265
A266
K274
L277
E280
C281
N282
T296
P315
D322
L328
H334
S335
A336
GLN

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	67.18Å 83.01Å 277.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.88 – 2.00 28.73 – 2.00	Depositor EDS
% Data completeness (in resolution range)	89.9 (28.88-2.00) 89.9 (28.73-2.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.58 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, R_{free}	0.213 , 0.266 0.213 , 0.266	Depositor DCC
R_{free} test set	4767 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	21.3	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 63.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.42$, $< L^2 > = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	10895	wwPDB-VP
Average B, all atoms (Å^2)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: R15

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.59	0/2485	0.68	1/3356 (0.0%)
1	B	0.63	0/2469	0.69	1/3335 (0.0%)
1	C	0.63	0/2478	0.69	1/3346 (0.0%)
1	D	0.65	0/2469	0.68	0/3335
All	All	0.63	0/9901	0.69	3/13372 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	33	LEU	CA-CB-CG	6.91	131.20	115.30
1	A	278	LEU	CA-CB-CG	5.91	128.88	115.30
1	C	254	ARG	NE-CZ-NH2	-5.52	117.54	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2442	0	2491	43	0
1	B	2426	0	2469	40	0
1	C	2435	0	2482	34	0
1	D	2426	0	2469	47	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	42	0	32	19	0
2	D	42	0	32	21	0
3	A	217	0	0	0	0
3	B	270	0	0	1	0
3	C	292	0	0	7	0
3	D	303	0	0	7	0
All	All	10895	0	9975	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:MET:CE	2:A:1336[A]:R15:H512	1.38	1.53
1:A:18:MET:CE	2:A:1336[A]:R15:C51	1.92	1.46
1:A:18:MET:HE1	2:A:1336[A]:R15:C51	1.59	1.23
1:A:18:MET:HE3	2:A:1336[A]:R15:C51	1.56	1.18
1:D:18:MET:CE	2:D:1336[A]:R15:H512	1.76	1.15

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	316/338 (94%)	305 (96%)	11 (4%)	0	100 100
1	B	314/338 (93%)	303 (96%)	11 (4%)	0	100 100
1	C	315/338 (93%)	301 (96%)	13 (4%)	1 (0%)	41 37
1	D	314/338 (93%)	305 (97%)	9 (3%)	0	100 100
All	All	1259/1352 (93%)	1214 (96%)	44 (4%)	1 (0%)	51 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	335	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/281 (95%)	259 (97%)	7 (3%)	46	48
1	B	264/281 (94%)	255 (97%)	9 (3%)	37	36
1	C	265/281 (94%)	262 (99%)	3 (1%)	73	78
1	D	264/281 (94%)	258 (98%)	6 (2%)	50	53
All	All	1059/1124 (94%)	1034 (98%)	25 (2%)	49	51

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	282	ASN
1	C	200	VAL
1	D	335	SER
1	C	56	LEU
1	C	264	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	182	ASN
1	D	334	HIS
1	C	182	ASN
1	C	282	ASN
1	C	334	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	R15	A	1336[B]	-	36,36,36	1.35	5 (13%)	49,49,49	1.29	4 (8%)
2	R15	D	1336[A]	-	36,36,36	1.41	6 (16%)	49,49,49	1.89	12 (24%)
2	R15	A	1336[A]	-	36,36,36	1.35	5 (13%)	49,49,49	1.27	4 (8%)
2	R15	D	1336[B]	-	36,36,36	1.41	6 (16%)	49,49,49	1.89	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R15	A	1336[B]	-	-	3/34/34/34	0/2/2/2
2	R15	D	1336[A]	-	-	4/34/34/34	0/2/2/2
2	R15	A	1336[A]	-	-	4/34/34/34	0/2/2/2
2	R15	D	1336[B]	-	-	3/34/34/34	0/2/2/2

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1336[A]	R15	O14-S1	4.49	1.48	1.43
2	A	1336[B]	R15	O14-S1	4.49	1.48	1.43
2	D	1336[A]	R15	O11-S1	3.99	1.48	1.43
2	D	1336[B]	R15	O11-S1	3.99	1.48	1.43
2	D	1336[A]	R15	O14-S1	3.61	1.47	1.43

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	D	1336[A]	R15	O12-S2-C10	5.37	114.58	107.97
2	D	1336[B]	R15	O12-S2-C10	5.37	114.58	107.97
2	D	1336[A]	R15	O14-S1-O11	-5.09	113.29	119.55
2	D	1336[B]	R15	O14-S1-O11	-5.09	113.29	119.55
2	A	1336[A]	R15	O14-S1-O11	-4.75	113.71	119.55

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1336[B]	R15	C35-C45-C51-C57
2	D	1336[B]	R15	C35-C45-C51-C57
2	A	1336[B]	R15	N21-C38-C48-C54
2	A	1336[A]	R15	N22-C35-C45-C51
2	D	1336[B]	R15	C48-C54-C57-C51

There are no ring outliers.

4 monomers are involved in 40 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1336[B]	R15	7	0
2	D	1336[A]	R15	12	0
2	A	1336[A]	R15	12	0
2	D	1336[B]	R15	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand R15 A 1336 (B)



Ligand R15 D 1336 (A)



Ligand R15 A 1336 (A)





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	320/338 (94%)	-0.04	5 (1%)	72	70	11, 23, 40, 46	0
1	B	318/338 (94%)	-0.11	9 (2%)	53	51	11, 21, 37, 46	0
1	C	319/338 (94%)	-0.19	3 (0%)	84	83	12, 21, 35, 45	0
1	D	318/338 (94%)	-0.21	2 (0%)	89	88	10, 19, 34, 46	0
All	All	1275/1352 (94%)	-0.14	19 (1%)	73	72	10, 21, 37, 46	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	336	ALA	6.8
1	A	336	ALA	4.1
1	B	336	ALA	3.8
1	B	142	LYS	3.2
1	D	236	ASN	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	R15	A	1336[A]	35/35	0.96	0.12	29,31,34,35	7
2	R15	A	1336[B]	35/35	0.96	0.12	29,31,33,35	7
2	R15	D	1336[A]	35/35	0.96	0.12	28,31,34,34	7
2	R15	D	1336[B]	35/35	0.96	0.12	28,30,33,34	7

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around R15 A 1336 (A):

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around R15 A 1336 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around R15 D 1336 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around R15 D 1336 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.