



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 04:29 PM EDT

PDB ID : 3LVF
Title : Crystal Structure of holo Glyceraldehyde-3-phosphate dehydrogenase 1 (GAPDH1) from methicillin resistant Staphylococcus aureus MRSA252 at 1.7 Angstrom resolution
Authors : Mukherjee, S.; Dutta, D.; Saha, B.; Das, A.K.
Deposited on : 2010-02-19
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	338	 6% 93% 5% ..
1	P	338	 9% 95% 5%
1	Q	338	 18% 95% ..
1	R	338	 9% 93% 6% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	P	338	2575	1603	441	521	10	0	4	0
1	R	335	2566	1597	441	519	9	0	5	0
1	O	334	2535	1578	435	513	9	0	2	0
1	Q	334	2529	1574	435	511	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	-1	GLY	-	expression tag	UNP Q6GIL8
P	0	SER	-	expression tag	UNP Q6GIL8
R	-1	GLY	-	expression tag	UNP Q6GIL8
R	0	SER	-	expression tag	UNP Q6GIL8
O	-1	GLY	-	expression tag	UNP Q6GIL8
O	0	SER	-	expression tag	UNP Q6GIL8
Q	-1	GLY	-	expression tag	UNP Q6GIL8
Q	0	SER	-	expression tag	UNP Q6GIL8

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	P	1	44	21	7	14	2	0	0
2	R	1	44	21	7	14	2	0	0
2	O	1	44	21	7	14	2	0	0
2	Q	1	44	21	7	14	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	168	Total	O	0	0
			168	168		
3	R	184	Total	O	0	0
			184	184		
3	O	181	Total	O	0	0
			181	181		
3	Q	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase 1



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase 1




- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase 1



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.18Å 104.87Å 91.19Å 90.00° 107.72° 90.00°	Depositor
Resolution (Å)	20.00 – 1.70 30.78 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.5 (20.00-1.70) 94.5 (30.78-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0095	Depositor
R, R_{free}	0.189 , 0.221 0.200 , 0.232	Depositor DCC
R_{free} test set	6359 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtrriage
Anisotropy	0.198	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11052	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	O	0.35	0/2575	0.50	0/3489
1	P	0.35	0/2621	0.51	0/3548
1	Q	0.34	0/2563	0.50	0/3473
1	R	0.35	0/2612	0.51	0/3538
All	All	0.35	0/10371	0.51	0/14048

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	2535	0	2515	17	0
1	P	2575	0	2564	9	0
1	Q	2529	0	2505	11	0
1	R	2566	0	2550	15	0
2	O	44	0	26	0	0
2	P	44	0	26	0	0
2	Q	44	0	26	1	0
2	R	44	0	26	0	0
3	O	181	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	P	168	0	0	2	0
3	Q	138	0	0	0	0
3	R	184	0	0	1	0
All	All	11052	0	10238	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:53:ARG:HH11	1:O:53:ARG:HG3	1.40	0.84
1:R:105:LYS:HE2	3:R:582:HOH:O	1.84	0.75
3:P:450:HOH:O	1:O:298:ARG:HD3	1.87	0.74
1:R:3:VAL:HG21	1:R:332:ALA:HB1	1.79	0.63
1:R:131:VAL:H	1:R:135:ASN:HD21	1.47	0.61
1:O:183:ASP:O	1:O:191:HIS:HE1	1.84	0.59
1:R:131:VAL:H	1:R:135:ASN:ND2	2.01	0.59
1:O:3:VAL:HG21	1:O:332:ALA:HB1	1.89	0.54
1:O:53:ARG:HG3	1:O:53:ARG:NH1	2.15	0.51
1:O:315:ASP:O	1:O:319:SER:HB2	2.10	0.51
1:P:155:SER:HA	1:P:290:SER:HB2	1.92	0.51
1:R:173:LEU:HD13	1:Q:309:LYS:HB2	1.92	0.51
1:Q:83:LEU:HD13	1:Q:85:TRP:CZ2	2.45	0.50
1:Q:211:THR:HG22	1:Q:231:GLY:HA2	1.94	0.50
1:O:191:HIS:CD2	1:O:193:LYS:H	2.30	0.49
1:P:18:PHE:O	1:P:22:GLN:HG2	2.14	0.48
1:O:83:LEU:HD13	1:O:85:TRP:CZ2	2.49	0.47
1:R:155:SER:HA	1:R:290:SER:HB2	1.97	0.47
1:R:211:THR:HG22	1:R:231:GLY:HA2	1.95	0.47
1:O:155:SER:HA	1:O:290:SER:HB2	1.97	0.46
1:O:268:ASN:ND2	1:O:271:PHE:H	2.14	0.46
1:R:130:ILE:HA	1:R:135:ASN:HD21	1.81	0.46
1:R:83:LEU:HD13	1:R:85:TRP:CZ2	2.51	0.45
1:R:99:PHE:O	1:R:105:LYS:HE3	2.17	0.45
1:Q:155:SER:HA	1:Q:290:SER:HB2	1.99	0.45
1:P:205:ASN:HD21	1:O:281:SER:CB	2.31	0.44
1:Q:85:TRP:HA	1:Q:85:TRP:CE3	2.53	0.44
1:R:205:ASN:HD21	1:Q:281:SER:CB	2.31	0.43
1:O:268:ASN:HD22	1:O:270:SER:H	1.65	0.43
1:O:85:TRP:HA	1:O:85:TRP:CE3	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:85:TRP:HA	1:O:85:TRP:HE3	1.83	0.43
1:R:281:SER:CB	1:Q:205:ASN:HD21	2.31	0.43
1:O:268:ASN:ND2	1:O:270:SER:H	2.17	0.43
1:P:263:MET:HG3	1:P:292:PHE:CZ	2.54	0.43
1:Q:85:TRP:HA	1:Q:85:TRP:HE3	1.83	0.43
1:P:83:LEU:HD13	1:P:85:TRP:CZ2	2.54	0.43
1:R:85:TRP:CE3	1:R:85:TRP:HA	2.54	0.43
1:O:268:ASN:HD22	1:O:268:ASN:C	2.22	0.42
1:R:85:TRP:HA	1:R:85:TRP:HE3	1.84	0.42
1:R:3:VAL:HB	1:R:27:LEU:HD12	2.00	0.42
1:Q:315:ASP:O	1:Q:319:SER:HB2	2.19	0.42
1:P:6:ALA:HB2	1:P:90:ILE:HD12	2.01	0.42
1:Q:11:GLY:HA3	2:Q:337:NAD:O5B	2.19	0.42
1:Q:34:ASP:O	1:Q:76:SER:HA	2.20	0.42
1:O:53:ARG:NH1	1:O:53:ARG:CG	2.81	0.41
1:P:61:VAL:HG11	1:P:66:ARG:HH11	1.86	0.41
1:P:211:THR:HG22	1:P:231:GLY:HA2	2.01	0.41
1:P:234:ARG:HD2	3:P:689:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	334/338 (99%)	322 (96%)	12 (4%)	0	100	100
1	P	340/338 (101%)	327 (96%)	13 (4%)	0	100	100
1	Q	332/338 (98%)	318 (96%)	14 (4%)	0	100	100
1	R	338/338 (100%)	325 (96%)	13 (4%)	0	100	100
All	All	1344/1352 (99%)	1292 (96%)	52 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	274/275 (100%)	269 (98%)	5 (2%)	59	43
1	P	279/275 (102%)	277 (99%)	2 (1%)	84	77
1	Q	272/275 (99%)	271 (100%)	1 (0%)	91	87
1	R	278/275 (101%)	272 (98%)	6 (2%)	52	34
All	All	1103/1100 (100%)	1089 (99%)	14 (1%)	69	56

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	85	TRP
1	P	209	ASN
1	R	85	TRP
1	R	127	LEU
1	R	135	ASN
1	R	209	ASN
1	R	300	MET
1	R	307	LEU
1	O	71	GLU
1	O	85	TRP
1	O	193	LYS
1	O	209	ASN
1	O	268	ASN
1	Q	85	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	P	68	ASN
1	P	205	ASN
1	P	252	GLN
1	R	68	ASN
1	R	135	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	R	205	ASN
1	R	265	ASN
1	R	296	GLN
1	O	68	ASN
1	O	191	HIS
1	O	205	ASN
1	O	265	ASN
1	O	268	ASN
1	O	296	GLN
1	Q	68	ASN
1	Q	205	ASN
1	Q	265	ASN
1	Q	296	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	P	337	-	42,48,48	1.73	4 (9%)	50,73,73	1.17	2 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	Q	337	-	42,48,48	1.76	4 (9%)	50,73,73	1.16	2 (4%)
2	NAD	O	337	-	42,48,48	1.74	4 (9%)	50,73,73	1.20	2 (4%)
2	NAD	R	337	-	42,48,48	1.74	4 (9%)	50,73,73	1.18	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	P	337	-	-	6/26/62/62	0/5/5/5
2	NAD	Q	337	-	-	5/26/62/62	0/5/5/5
2	NAD	O	337	-	-	5/26/62/62	0/5/5/5
2	NAD	R	337	-	-	5/26/62/62	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Q	337	NAD	O7N-C7N	8.91	1.41	1.24
2	R	337	NAD	O7N-C7N	8.85	1.41	1.24
2	O	337	NAD	O7N-C7N	8.82	1.41	1.24
2	P	337	NAD	O7N-C7N	8.70	1.40	1.24
2	Q	337	NAD	C2A-N3A	4.19	1.38	1.32
2	R	337	NAD	C2A-N3A	4.11	1.38	1.32
2	O	337	NAD	C2A-N3A	4.08	1.38	1.32
2	P	337	NAD	C2A-N3A	4.04	1.38	1.32
2	O	337	NAD	C2N-N1N	2.61	1.38	1.35
2	Q	337	NAD	C2A-N1A	2.56	1.38	1.33
2	P	337	NAD	C2N-N1N	2.53	1.38	1.35
2	Q	337	NAD	C2N-N1N	2.51	1.38	1.35
2	R	337	NAD	C2A-N1A	2.51	1.38	1.33
2	P	337	NAD	C2A-N1A	2.45	1.38	1.33
2	O	337	NAD	C2A-N1A	2.42	1.38	1.33
2	R	337	NAD	C2N-N1N	2.18	1.37	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	O	337	NAD	N3A-C2A-N1A	-5.92	119.42	128.68
2	R	337	NAD	N3A-C2A-N1A	-5.78	119.64	128.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	337	NAD	N3A-C2A-N1A	-5.73	119.72	128.68
2	Q	337	NAD	N3A-C2A-N1A	-5.68	119.80	128.68
2	Q	337	NAD	PN-O3-PA	-2.53	124.13	132.83
2	O	337	NAD	PN-O3-PA	-2.38	124.66	132.83
2	P	337	NAD	PN-O3-PA	-2.32	124.88	132.83
2	R	337	NAD	PN-O3-PA	-2.20	125.29	132.83
2	R	337	NAD	C1B-N9A-C4A	-2.17	122.83	126.64

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	P	337	NAD	O4D-C1D-N1N-C2N
2	P	337	NAD	O4D-C1D-N1N-C6N
2	P	337	NAD	C2D-C1D-N1N-C2N
2	P	337	NAD	C2D-C1D-N1N-C6N
2	R	337	NAD	O4D-C1D-N1N-C2N
2	R	337	NAD	O4D-C1D-N1N-C6N
2	R	337	NAD	C2D-C1D-N1N-C6N
2	O	337	NAD	O4D-C1D-N1N-C2N
2	O	337	NAD	O4D-C1D-N1N-C6N
2	O	337	NAD	C2D-C1D-N1N-C6N
2	Q	337	NAD	O4D-C1D-N1N-C2N
2	Q	337	NAD	O4D-C1D-N1N-C6N
2	Q	337	NAD	C2D-C1D-N1N-C6N
2	P	337	NAD	O4B-C4B-C5B-O5B
2	O	337	NAD	O4B-C4B-C5B-O5B
2	Q	337	NAD	O4B-C4B-C5B-O5B
2	R	337	NAD	O4B-C4B-C5B-O5B
2	P	337	NAD	C3B-C4B-C5B-O5B
2	R	337	NAD	C2D-C1D-N1N-C2N
2	O	337	NAD	C2D-C1D-N1N-C2N
2	Q	337	NAD	C2D-C1D-N1N-C2N


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Q	337	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	O	334/338 (98%)	0.40	19 (5%) 23 26	29, 36, 47, 56	3 (0%)
1	P	338/338 (100%)	0.63	29 (8%) 10 12	29, 37, 50, 59	3 (0%)
1	Q	334/338 (98%)	1.05	62 (18%) 1 1	29, 40, 61, 68	0
1	R	335/338 (99%)	0.63	30 (8%) 9 10	29, 36, 48, 64	5 (1%)
All	All	1341/1352 (99%)	0.68	140 (10%) 6 7	29, 37, 54, 68	11 (0%)

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Q	69	GLY	9.4
1	R	335	SER	9.2
1	Q	2	ALA	6.8
1	O	55	THR	6.6
1	P	334	LEU	5.8
1	P	55	THR	5.6
1	R	334	LEU	5.5
1	Q	24	VAL	5.5
1	Q	304	ASP	5.4
1	O	69	GLY	5.3
1	P	192	ARG	5.2
1	Q	99	PHE	5.1
1	Q	63	GLY	5.1
1	P	-1	GLY	5.0
1	R	55	THR	4.9
1	P	62[A]	ASP	4.9
1	R	126	ASP	4.8
1	P	1	MET	4.8
1	O	334	LEU	4.8
1	R	333	GLU	4.7
1	Q	88	LEU	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	R	336	LYS	4.5
1	R	66	ARG	4.4
1	Q	89	ASN	4.3
1	R	304	ASP	4.2
1	P	126[A]	ASP	4.2
1	Q	13	ILE	4.2
1	P	56	GLY	4.2
1	Q	192	ARG	4.2
1	Q	62	ASP	4.1
1	Q	61	VAL	3.9
1	Q	334	LEU	3.9
1	P	333	GLU	3.9
1	Q	25	GLU	3.7
1	R	324	LEU	3.7
1	P	66	ARG	3.6
1	O	62	ASP	3.6
1	O	66	ARG	3.6
1	Q	60	VAL	3.6
1	R	39[A]	ASP	3.5
1	R	192	ARG	3.5
1	Q	3	VAL	3.5
1	Q	86	LYS	3.5
1	Q	68	ASN	3.5
1	Q	125	GLY	3.4
1	R	68	ASN	3.3
1	Q	332	ALA	3.3
1	Q	71	GLU	3.3
1	P	156	LEU	3.3
1	R	69	GLY	3.3
1	Q	107	GLN	3.3
1	R	156	LEU	3.3
1	Q	70	LYS	3.2
1	Q	261	GLU	3.2
1	Q	333	GLU	3.2
1	O	335	SER	3.2
1	Q	242	LEU	3.2
1	Q	252	GLN	3.2
1	Q	102	ASP	3.1
1	P	57	GLU	3.1
1	Q	87	ASP	3.1
1	P	142	SER	3.1
1	Q	55	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	R	328	LEU	3.0
1	R	57	GLU	3.0
1	R	141	GLY	3.0
1	Q	124	THR	2.9
1	R	140	ASP	2.9
1	O	57	GLU	2.9
1	Q	137	GLN	2.9
1	P	252	GLN	2.9
1	Q	64	GLY	2.9
1	P	71	GLU	2.9
1	R	320	TYR	2.9
1	P	304	ASP	2.9
1	O	61	VAL	2.9
1	P	140	ASP	2.8
1	Q	126	ASP	2.8
1	Q	53	ARG	2.8
1	Q	39	ASP	2.8
1	Q	23	GLU	2.7
1	P	69	GLY	2.7
1	Q	156	LEU	2.6
1	P	70	LYS	2.6
1	Q	90	ILE	2.6
1	Q	72	VAL	2.6
1	R	325	VAL	2.5
1	Q	320	TYR	2.5
1	Q	103	LYS	2.5
1	O	68	ASN	2.5
1	Q	36	THR	2.5
1	O	142	SER	2.5
1	P	13	ILE	2.5
1	P	141	GLY	2.5
1	P	242	LEU	2.4
1	Q	104	ASP	2.4
1	Q	335	SER	2.4
1	O	56	GLY	2.4
1	R	16	LEU	2.4
1	Q	253	ASP	2.4
1	Q	121	ALA	2.4
1	P	84	PRO	2.3
1	R	13	ILE	2.3
1	Q	27	LEU	2.3
1	R	332	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	O	137	GLN	2.3
1	Q	76	SER	2.3
1	O	192	ARG	2.3
1	Q	28	GLU	2.3
1	P	335	SER	2.3
1	Q	251	LYS	2.3
1	P	177	ILE	2.3
1	O	59	GLU	2.3
1	O	333	GLU	2.2
1	P	151	CYS	2.2
1	O	28	GLU	2.2
1	O	25	GLU	2.1
1	R	7	ILE	2.1
1	O	177	ILE	2.1
1	R	310	VAL	2.1
1	Q	29	VAL	2.1
1	Q	96	CYS	2.1
1	P	0	SER	2.1
1	R	94	LEU	2.1
1	R	59	GLU	2.1
1	P	253	ASP	2.1
1	Q	7	ILE	2.1
1	Q	108	ALA	2.0
1	O	99	PHE	2.0
1	Q	10	PHE	2.0
1	R	291	LEU	2.0
1	R	321	THR	2.0
1	Q	37	ASP	2.0
1	R	2	ALA	2.0
1	Q	193	LYS	2.0
1	Q	314	TYR	2.0
1	Q	65	PHE	2.0
1	Q	118	LEU	2.0
1	P	89	ASN	2.0
1	Q	141	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAD	Q	337	44/44	0.92	0.10	32,35,38,38	0
2	NAD	R	337	44/44	0.93	0.10	31,34,36,37	0
2	NAD	P	337	44/44	0.93	0.10	29,34,35,36	0
2	NAD	O	337	44/44	0.94	0.10	31,34,36,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAD R 337:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD P 337:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.