



# Full wwPDB X-ray Structure Validation Report i

May 22, 2020 – 08:27 pm BST

PDB ID : 5MBQ  
Title : CeuE (H227A variant) a periplasmic protein from *Campylobacter jejuni*  
Authors : Wilde, E.J.; Blagova, E.V.; Hughes, A.; Raines, D.J.; Moroz, O.V.; Turkenburg, J.P.; Duhme-Klair, A.-K.; Wilson, K.S.  
Deposited on : 2016-11-08  
Resolution : 1.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 7062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enterochelin uptake periplasmic binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	286	Total	C 2189	N 1417	O 347	S 422	3	0	10	0
1	B	287	Total	C 2194	N 1423	O 346	S 422	3	0	11	1
1	C	282	Total	C 2156	N 1399	O 344	S 410	3	0	5	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLY	-	expression tag	UNP Q0P8Q4
A	21	PRO	-	expression tag	UNP Q0P8Q4
A	22	ALA	-	expression tag	UNP Q0P8Q4
A	23	MET	-	expression tag	UNP Q0P8Q4
A	227	ALA	HIS	engineered mutation	UNP Q0P8Q4
B	20	GLY	-	expression tag	UNP Q0P8Q4
B	21	PRO	-	expression tag	UNP Q0P8Q4
B	22	ALA	-	expression tag	UNP Q0P8Q4
B	23	MET	-	expression tag	UNP Q0P8Q4
B	227	ALA	HIS	engineered mutation	UNP Q0P8Q4
C	20	GLY	-	expression tag	UNP Q0P8Q4
C	21	PRO	-	expression tag	UNP Q0P8Q4
C	22	ALA	-	expression tag	UNP Q0P8Q4
C	23	MET	-	expression tag	UNP Q0P8Q4
C	227	ALA	HIS	engineered mutation	UNP Q0P8Q4

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	199	Total 199 199	0	0
2	B	181	Total 181 181	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	143	Total O 143 143	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enterochelin uptake periplasmic binding protein



- Molecule 1: Enterochelin uptake periplasmic binding protein



- Molecule 1: Enterochelin uptake periplasmic binding protein



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.99 Å   62.68 Å   67.97 Å 82.11°   77.21°   76.28°	Depositor
Resolution (Å)	66.02 – 1.33 66.02 – 1.33	Depositor EDS
% Data completeness (in resolution range)	87.2 (66.02-1.33) 87.2 (66.02-1.33)	Depositor EDS
$R_{merge}$	0.00	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.11 (at 1.33 Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
$R$ , $R_{free}$	0.148 , 0.193 0.153 , 0.201	Depositor DCC
$R_{free}$ test set	8932 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.5	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.4	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.47$ , $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	7062	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.69	19/2250 (0.8%)	1.44	22/3045 (0.7%)
1	B	1.57	13/2256 (0.6%)	1.38	17/3049 (0.6%)
1	C	1.63	21/2202 (1.0%)	1.52	18/2984 (0.6%)
All	All	1.63	53/6708 (0.8%)	1.45	57/9078 (0.6%)

All (53) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	116	SER	CB-OG	12.50	1.58	1.42
1	A	220	GLU	CD-OE2	8.72	1.35	1.25
1	A	286	TYR	CE1-CZ	-7.83	1.28	1.38
1	B	128[A]	GLU	CD-OE1	7.59	1.34	1.25
1	B	128[B]	GLU	CD-OE1	7.59	1.34	1.25
1	B	296	GLU	CG-CD	7.47	1.63	1.51
1	C	145	SER	CB-OG	-7.25	1.32	1.42
1	A	285	GLU	CD-OE2	-7.25	1.17	1.25
1	A	175	GLU	CD-OE1	-7.16	1.17	1.25
1	A	286	TYR	CD1-CE1	7.15	1.50	1.39
1	B	234	GLU	CD-OE1	7.10	1.33	1.25
1	B	59	GLY	CA-C	-7.09	1.40	1.51
1	C	228	GLY	N-CA	6.85	1.56	1.46
1	B	228	GLY	N-CA	6.82	1.56	1.46
1	C	157	TYR	CE1-CZ	-6.79	1.29	1.38
1	A	294	GLY	N-CA	6.58	1.55	1.46
1	B	285	GLU	CD-OE1	6.28	1.32	1.25
1	B	179	SER	CB-OG	-6.17	1.34	1.42
1	A	71	ASP	CB-CG	6.14	1.64	1.51
1	B	243	TYR	CG-CD1	-6.10	1.31	1.39
1	B	175	GLU	CD-OE2	6.09	1.32	1.25
1	C	111	ASP	N-CA	-6.08	1.34	1.46
1	C	157	TYR	CG-CD2	-6.07	1.31	1.39
1	C	64	PHE	CE1-CZ	5.94	1.48	1.37
1	A	157	TYR	CE2-CZ	5.83	1.46	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	88	PHE	CE1-CZ	5.81	1.48	1.37
1	A	286	TYR	CZ-OH	5.80	1.47	1.37
1	C	281	TYR	CG-CD2	5.77	1.46	1.39
1	C	69	LEU	C-O	5.66	1.34	1.23
1	B	135	VAL	C-O	-5.64	1.12	1.23
1	B	40	SER	CB-OG	-5.61	1.34	1.42
1	A	60	ILE	CA-C	5.54	1.67	1.52
1	C	296	GLU	CD-OE2	5.53	1.31	1.25
1	C	118	ARG	CZ-NH1	5.47	1.40	1.33
1	C	128	GLU	CD-OE2	5.40	1.31	1.25
1	A	286	TYR	CD2-CE2	5.35	1.47	1.39
1	A	110	PRO	N-CA	-5.34	1.38	1.47
1	C	84	TYR	CE1-CZ	5.33	1.45	1.38
1	C	249	ARG	N-CA	5.26	1.56	1.46
1	C	173	GLU	CG-CD	5.25	1.59	1.51
1	C	193	ASN	CB-CG	-5.21	1.39	1.51
1	C	147	GLU	CD-OE2	5.16	1.31	1.25
1	A	243	TYR	CG-CD1	-5.14	1.32	1.39
1	C	49[A]	ASN	CG-OD1	5.12	1.35	1.24
1	C	49[B]	ASN	CG-OD1	5.12	1.35	1.24
1	A	271	THR	CB-OG1	5.11	1.53	1.43
1	A	152	SER	N-CA	-5.08	1.36	1.46
1	A	64	PHE	CG-CD1	5.08	1.46	1.38
1	B	258	ARG	CZ-NH2	5.06	1.39	1.33
1	C	40	SER	CB-OG	-5.05	1.35	1.42
1	A	206	PHE	CG-CD1	5.04	1.46	1.38
1	A	146	PHE	N-CA	5.02	1.56	1.46
1	A	142	PHE	CD2-CE2	5.00	1.49	1.39

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	28	MET	CG-SD-CE	-15.24	75.81	100.20
1	C	118	ARG	NE-CZ-NH1	13.59	127.10	120.30
1	C	118	ARG	NE-CZ-NH2	-12.62	113.99	120.30
1	A	28	MET	CG-SD-CE	-12.15	80.75	100.20
1	C	249	ARG	NE-CZ-NH2	-12.10	114.25	120.30
1	B	301	MET	CG-SD-CE	-11.83	81.27	100.20
1	A	283	ASP	CB-CG-OD2	-8.99	110.21	118.30
1	C	57	ASP	CB-CG-OD1	-8.34	110.79	118.30
1	C	81	LEU	CA-CB-CG	8.31	134.43	115.30
1	A	258	ARG	NE-CZ-NH2	-8.01	116.30	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	ASP	CB-CG-OD2	-7.91	111.18	118.30
1	A	101	ASP	CB-CG-OD1	7.86	125.37	118.30
1	B	84	TYR	CD1-CE1-CZ	-7.32	113.21	119.80
1	C	180	ILE	CG1-CB-CG2	7.11	127.05	111.40
1	C	24	LEU	CA-CB-CG	6.91	131.20	115.30
1	A	301	MET	CG-SD-CE	-6.85	89.23	100.20
1	C	283	ASP	CB-CG-OD2	-6.66	112.31	118.30
1	B	287	TRP	CD1-NE1-CE2	6.63	114.97	109.00
1	A	286	TYR	CG-CD1-CE1	-6.54	116.07	121.30
1	C	23	MET	CG-SD-CE	-6.44	89.90	100.20
1	C	64	PHE	CB-CG-CD1	6.40	125.28	120.80
1	B	249	ARG	NE-CZ-NH2	-6.35	117.13	120.30
1	A	175	GLU	OE1-CD-OE2	-6.27	115.78	123.30
1	B	281	TYR	CB-CG-CD2	6.21	124.73	121.00
1	B	258	ARG	NE-CZ-NH1	-6.11	117.25	120.30
1	B	286	TYR	CB-CG-CD2	-6.11	117.33	121.00
1	C	157	TYR	CB-CG-CD1	-6.10	117.34	121.00
1	A	65	ASP	CB-CG-OD1	6.09	123.78	118.30
1	A	249	ARG	NE-CZ-NH2	6.05	123.32	120.30
1	A	146	PHE	CB-CG-CD1	-6.02	116.58	120.80
1	A	81	LEU	CB-CG-CD1	6.02	121.23	111.00
1	C	301	MET	CG-SD-CE	-5.99	90.62	100.20
1	C	249	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	B	125	LYS	CD-CE-NZ	5.94	125.36	111.70
1	A	39	ASP	CB-CG-OD1	5.94	123.64	118.30
1	C	64	PHE	CB-CG-CD2	-5.93	116.65	120.80
1	C	182	ASP	CB-CG-OD1	5.90	123.61	118.30
1	A	286	TYR	CB-CG-CD2	-5.86	117.49	121.00
1	B	28	MET	CG-SD-CE	-5.83	90.88	100.20
1	A	211	ASP	CB-CG-OD2	5.81	123.53	118.30
1	B	205	ARG	NE-CZ-NH1	-5.80	117.40	120.30
1	B	23	MET	CA-CB-CG	5.77	123.11	113.30
1	C	296	GLU	OE1-CD-OE2	5.66	130.09	123.30
1	A	157	TYR	CB-CG-CD2	5.64	124.39	121.00
1	B	169	ASP	CB-CG-OD2	-5.64	113.22	118.30
1	A	142	PHE	CB-CG-CD1	-5.54	116.92	120.80
1	A	283	ASP	OD1-CG-OD2	5.44	133.64	123.30
1	A	286	TYR	CD1-CG-CD2	5.40	123.84	117.90
1	B	219	ASP	CB-CG-OD2	-5.37	113.47	118.30
1	A	63	THR	O-C-N	5.26	131.12	122.70
1	A	157	TYR	CB-CG-CD1	-5.25	117.85	121.00
1	A	287	TRP	CD1-CG-CD2	-5.23	102.11	106.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	ASP	CB-CG-OD1	-5.19	113.63	118.30
1	A	102	PHE	CB-CG-CD1	-5.18	117.17	120.80
1	C	164	LEU	CB-CG-CD2	5.15	119.76	111.00
1	B	229	LYS	O-C-N	-5.03	114.66	122.70
1	B	287	TRP	NE1-CE2-CD2	-5.01	102.29	107.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2189	0	2209	19	0
1	B	2194	0	2211	20	1
1	C	2156	0	2166	19	1
2	A	199	0	0	6	1
2	B	181	0	0	3	0
2	C	143	0	0	3	1
All	All	7062	0	6586	58	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:MET:SD	1:A:23:MET:CE	2.02	1.47
1:B:195:ASN:HB2	2:B:495:HOH:O	1.42	1.20
1:C:234:GLU:CG	2:C:530:HOH:O	1.91	1.16
1:A:26:ILE:H	1:A:148:ASN:HD21	1.20	0.88
1:B:26:ILE:H	1:B:148:ASN:HD21	1.23	0.86
1:C:220:GLU:N	1:C:220:GLU:OE1	2.10	0.85
1:C:26:ILE:H	1:C:148:ASN:HD21	1.27	0.82
1:A:96:GLY:O	2:A:402:HOH:O	1.96	0.81
1:B:220:GLU:CB	2:B:565:HOH:O	2.29	0.80

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83:LYS:CG	2:C:513:HOH:O	2.31	0.78
1:A:70:ASN:H	1:A:70:ASN:HD22	1.38	0.71
1:A:22:ALA:N	2:A:403:HOH:O	2.25	0.70
1:C:78:ALA:HB3	1:C:93:SER:HB2	1.75	0.69
1:A:193:ASN:HB3	2:A:411:HOH:O	1.94	0.67
1:C:250:ASN:HD22	1:C:255:ASN:HD22	1.43	0.67
1:B:250:ASN:HD22	1:B:255:ASN:HD22	1.51	0.58
1:A:121:LYS:CE	2:A:425:HOH:O	2.51	0.58
1:B:70:ASN:H	1:B:70:ASN:HD22	1.52	0.58
1:B:26:ILE:N	1:B:148:ASN:HD21	1.98	0.56
1:B:210:HIS:HD2	1:B:215:ILE:O	1.89	0.55
1:C:219:ASP:C	1:C:220:GLU:OE1	2.45	0.55
1:C:210:HIS:HD2	1:C:215:ILE:O	1.90	0.55
1:C:70[A]:ASN:HD22	1:C:70[A]:ASN:H	1.55	0.55
1:A:170[A]:ILE:HD11	1:A:295:LEU:O	2.07	0.54
1:A:23:MET:CG	1:A:23:MET:CE	2.85	0.54
1:C:65:ASP:HA	1:C:70[B]:ASN:OD1	2.09	0.53
1:B:228:GLY:N	2:B:401:HOH:O	1.93	0.53
1:A:193:ASN:OD1	1:A:249:ARG:NH2	2.40	0.51
1:A:223:LYS:CA	2:A:564:HOH:O	2.58	0.51
1:A:210:HIS:HD2	1:A:215:ILE:O	1.95	0.50
1:B:26:ILE:H	1:B:148:ASN:ND2	2.02	0.50
1:C:194:SER:H	1:C:250:ASN:HD21	1.60	0.50
1:C:83:LYS:HA	1:C:86:GLN:HG2	1.94	0.49
1:C:250:ASN:ND2	1:C:255:ASN:HD22	2.10	0.48
1:A:70:ASN:H	1:A:70:ASN:ND2	2.11	0.47
1:A:210:HIS:HE1	2:A:446:HOH:O	1.97	0.47
1:C:249:ARG:NH1	1:C:253:LEU:HD11	2.29	0.46
1:C:208[B]:ILE:HG13	1:C:212:VAL:HB	1.96	0.46
1:B:194[B]:SER:H	1:B:250:ASN:HD21	1.64	0.45
1:C:210:HIS:HE1	2:C:422:HOH:O	1.98	0.45
1:A:115:ILE:HB	1:A:119:GLN:HB2	1.99	0.45
1:B:37:VAL:HG21	1:B:46:ILE:HD12	1.99	0.44
1:B:194[A]:SER:H	1:B:250:ASN:HD21	1.66	0.44
1:C:194:SER:N	1:C:250:ASN:HD21	2.14	0.43
1:B:174:ILE:HG13	1:B:302[B]:ILE:HD13	2.00	0.43
1:A:26:ILE:H	1:A:148:ASN:ND2	2.02	0.43
1:A:28:MET:HB3	1:A:28:MET:HE3	1.44	0.42
1:B:55:ILE:O	1:B:77:PRO:HD3	2.19	0.42
1:B:182:ASP:OD1	1:B:184:ASP:N	2.34	0.42
1:A:70:ASN:HD22	1:A:70:ASN:N	2.13	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:142:PHE:CG	1:C:208[A]:ILE:CG2	3.04	0.41
1:B:70:ASN:ND2	1:B:70:ASN:H	2.17	0.41
1:B:82:PRO:HB3	1:B:285:GLU:HB3	2.03	0.41
1:C:115:ILE:HB	1:C:119:GLN:HB2	2.02	0.41
1:A:258:ARG:HA	1:A:258:ARG:HD2	1.77	0.40
1:B:180:ILE:HD12	1:B:180:ILE:C	2.42	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:LYS:CG	2:C:523:HOH:O[1_456]	1.76	0.44
1:C:256:LYS:CG	2:A:570:HOH:O[1_565]	2.01	0.19

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	292/291 (100%)	283 (97%)	9 (3%)	0	100 100
1	B	293/291 (101%)	281 (96%)	12 (4%)	0	100 100
1	C	283/291 (97%)	272 (96%)	11 (4%)	0	100 100
All	All	868/873 (99%)	836 (96%)	32 (4%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	233/251 (93%)	227 (97%)	6 (3%)	46 12
1	B	233/251 (93%)	231 (99%)	2 (1%)	78 52
1	C	227/251 (90%)	221 (97%)	6 (3%)	46 12
All	All	693/753 (92%)	679 (98%)	14 (2%)	57 20

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	98	GLN
1	A	121	LYS
1	A	258	ARG
1	A	291[A]	SER
1	A	291[B]	SER
1	B	121	LYS
1	B	239	LYS
1	C	81	LEU
1	C	99	GLN
1	C	144	SER
1	C	165	GLU
1	C	180	ILE
1	C	191	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	90	ASN
1	A	148	ASN
1	A	172	ASN
1	A	210	HIS
1	A	276	ASN
1	B	70	ASN
1	B	148	ASN
1	B	210	HIS
1	B	250	ASN
1	C	90	ASN
1	C	98	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	148	ASN
1	C	210	HIS
1	C	216	ASN
1	C	250	ASN
1	C	307	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/291 (98%)	-0.11	7 (2%) 59 65	14, 23, 40, 49	5 (1%)
1	B	287/291 (98%)	-0.24	5 (1%) 70 74	16, 24, 38, 50	8 (2%)
1	C	282/291 (96%)	-0.19	1 (0%) 92 93	14, 25, 39, 62	3 (1%)
All	All	855/873 (97%)	-0.18	13 (1%) 73 77	14, 24, 39, 62	16 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	221	ASN	11.0
1	B	97	VAL	7.3
1	A	97	VAL	5.6
1	B	222	ILE	4.1
1	B	224	VAL	3.4
1	A	250	ASN	3.3
1	B	225	GLY	3.0
1	A	222	ILE	2.8
1	A	98	GLN	2.5
1	B	96	GLY	2.5
1	A	194	SER	2.4
1	A	258	ARG	2.3
1	A	224	VAL	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.