



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2023 – 12:32 pm GMT

PDB ID : 5MGS
Title : Human receptor NKR-P1 in deglycosylated form, extracellular domain
Authors : Skalova, T.; Blaha, J.; Stransky, J.; Koval, T.; Hasek, J.; Yuguang, Z.; Harlos, K.; Vanek, O.; Dohnalek, J.
Deposited on : 2016-11-22
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.32.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	81% (Poor fit: 0%, 3+ outliers: 0%, 2 outliers: 0%, 1 outlier: 0%, 0 outliers: 81%, Not modelled: 15%)
1	B	146	79% (Poor fit: 0%, 3+ outliers: 0%, 2 outliers: 0%, 1 outlier: 0%, 0 outliers: 79%, Not modelled: 12%)
1	C	146	2% (Poor fit: 2%, 3+ outliers: 0%, 2 outliers: 0%, 1 outlier: 0%, 0 outliers: 78%, Not modelled: 14%)
1	D	146	2% (Poor fit: 2%, 3+ outliers: 0%, 2 outliers: 0%, 1 outlier: 0%, 0 outliers: 81%, Not modelled: 14%)
1	E	146	2% (Poor fit: 2%, 3+ outliers: 0%, 2 outliers: 0%, 1 outlier: 0%, 0 outliers: 82%, Not modelled: 14%)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	146	<p>% 77% 8% 15%</p>
1	G	146	<p>% 82% 15%</p>
1	H	146	<p>% 84% 5% 12%</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Killer cell lectin-like receptor subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	124	1037	660	175	196	6	0	3	0
1	B	128	1053	670	179	198	6	0	1	0
1	C	126	1035	658	175	196	6	0	0	0
1	D	125	1030	654	175	195	6	0	1	0
1	E	125	1025	651	174	194	6	0	0	0
1	F	124	1037	660	175	196	6	0	4	0
1	G	124	1038	660	177	195	6	0	3	0
1	H	129	1075	682	186	201	6	0	3	0

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	GLU	-	expression tag	UNP Q12918
A	88	THR	-	expression tag	UNP Q12918
A	89	GLY	-	expression tag	UNP Q12918
A	226	LYS	-	expression tag	UNP Q12918
A	227	HIS	-	expression tag	UNP Q12918
A	228	HIS	-	expression tag	UNP Q12918
A	229	HIS	-	expression tag	UNP Q12918
A	230	HIS	-	expression tag	UNP Q12918
A	231	HIS	-	expression tag	UNP Q12918
A	232	HIS	-	expression tag	UNP Q12918
B	87	GLU	-	expression tag	UNP Q12918
B	88	THR	-	expression tag	UNP Q12918
B	89	GLY	-	expression tag	UNP Q12918

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	226	LYS	-	expression tag	UNP Q12918
B	227	HIS	-	expression tag	UNP Q12918
B	228	HIS	-	expression tag	UNP Q12918
B	229	HIS	-	expression tag	UNP Q12918
B	230	HIS	-	expression tag	UNP Q12918
B	231	HIS	-	expression tag	UNP Q12918
B	232	HIS	-	expression tag	UNP Q12918
C	87	GLU	-	expression tag	UNP Q12918
C	88	THR	-	expression tag	UNP Q12918
C	89	GLY	-	expression tag	UNP Q12918
C	226	LYS	-	expression tag	UNP Q12918
C	227	HIS	-	expression tag	UNP Q12918
C	228	HIS	-	expression tag	UNP Q12918
C	229	HIS	-	expression tag	UNP Q12918
C	230	HIS	-	expression tag	UNP Q12918
C	231	HIS	-	expression tag	UNP Q12918
C	232	HIS	-	expression tag	UNP Q12918
D	87	GLU	-	expression tag	UNP Q12918
D	88	THR	-	expression tag	UNP Q12918
D	89	GLY	-	expression tag	UNP Q12918
D	226	LYS	-	expression tag	UNP Q12918
D	227	HIS	-	expression tag	UNP Q12918
D	228	HIS	-	expression tag	UNP Q12918
D	229	HIS	-	expression tag	UNP Q12918
D	230	HIS	-	expression tag	UNP Q12918
D	231	HIS	-	expression tag	UNP Q12918
D	232	HIS	-	expression tag	UNP Q12918
E	87	GLU	-	expression tag	UNP Q12918
E	88	THR	-	expression tag	UNP Q12918
E	89	GLY	-	expression tag	UNP Q12918
E	226	LYS	-	expression tag	UNP Q12918
E	227	HIS	-	expression tag	UNP Q12918
E	228	HIS	-	expression tag	UNP Q12918
E	229	HIS	-	expression tag	UNP Q12918
E	230	HIS	-	expression tag	UNP Q12918
E	231	HIS	-	expression tag	UNP Q12918
E	232	HIS	-	expression tag	UNP Q12918
F	87	GLU	-	expression tag	UNP Q12918
F	88	THR	-	expression tag	UNP Q12918
F	89	GLY	-	expression tag	UNP Q12918
F	226	LYS	-	expression tag	UNP Q12918
F	227	HIS	-	expression tag	UNP Q12918

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	228	HIS	-	expression tag	UNP Q12918
F	229	HIS	-	expression tag	UNP Q12918
F	230	HIS	-	expression tag	UNP Q12918
F	231	HIS	-	expression tag	UNP Q12918
F	232	HIS	-	expression tag	UNP Q12918
G	87	GLU	-	expression tag	UNP Q12918
G	88	THR	-	expression tag	UNP Q12918
G	89	GLY	-	expression tag	UNP Q12918
G	226	LYS	-	expression tag	UNP Q12918
G	227	HIS	-	expression tag	UNP Q12918
G	228	HIS	-	expression tag	UNP Q12918
G	229	HIS	-	expression tag	UNP Q12918
G	230	HIS	-	expression tag	UNP Q12918
G	231	HIS	-	expression tag	UNP Q12918
G	232	HIS	-	expression tag	UNP Q12918
H	87	GLU	-	expression tag	UNP Q12918
H	88	THR	-	expression tag	UNP Q12918
H	89	GLY	-	expression tag	UNP Q12918
H	226	LYS	-	expression tag	UNP Q12918
H	227	HIS	-	expression tag	UNP Q12918
H	228	HIS	-	expression tag	UNP Q12918
H	229	HIS	-	expression tag	UNP Q12918
H	230	HIS	-	expression tag	UNP Q12918
H	231	HIS	-	expression tag	UNP Q12918
H	232	HIS	-	expression tag	UNP Q12918

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	C	1	14	8	1	5	0	0
2	D	1	14	8	1	5	0	0
2	E	1	14	8	1	5	0	0
2	E	1	14	8	1	5	0	0
2	F	1	14	8	1	5	0	0
2	F	1	14	8	1	5	0	0
2	G	1	14	8	1	5	0	0
2	G	1	14	8	1	5	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	H	1	Total	C	N	O	0	0
			14	8	1	5		
2	H	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	100	Total	O	0	0
			100	100		
3	B	92	Total	O	0	0
			92	92		
3	C	89	Total	O	0	0
			89	89		
3	D	72	Total	O	0	0
			72	72		
3	E	78	Total	O	0	0
			78	78		
3	F	86	Total	O	0	0
			86	86		
3	G	80	Total	O	0	0
			80	80		
3	H	85	Total	O	0	0
			85	85		



- Molecule 1: Killer cell lectin-like receptor subfamily B member 1



- Molecule 1: Killer cell lectin-like receptor subfamily B member 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	44.81Å 68.40Å 101.56Å 101.88° 100.72° 100.64°	Depositor
Resolution (Å)	48.68 – 1.90 48.68 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.9 (48.68-1.90) 97.9 (48.68-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.157 , 0.207 0.167 , 0.177	Depositor DCC
R_{free} test set	4289 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtrriage
Anisotropy	0.249	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9236	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	0/1069	0.93	2/1449 (0.1%)
1	B	0.89	0/1080	0.87	2/1465 (0.1%)
1	C	0.89	0/1058	0.86	3/1435 (0.2%)
1	D	0.88	0/1055	0.87	3/1429 (0.2%)
1	E	0.83	0/1047	0.81	0/1418
1	F	0.92	0/1073	0.88	1/1454 (0.1%)
1	G	0.89	0/1070	0.85	2/1450 (0.1%)
1	H	0.89	0/1107	0.87	1/1500 (0.1%)
All	All	0.89	0/8559	0.87	14/11600 (0.1%)

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	207	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	B	207	ARG	NE-CZ-NH2	7.00	123.80	120.30
1	C	146	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	F	102	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	H	121	ASP	CB-CG-OD2	6.59	124.23	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1037	0	1014	3	0
1	B	1053	0	1029	8	0
1	C	1035	0	1009	6	0
1	D	1030	0	1006	10	0
1	E	1025	0	999	5	0
1	F	1037	0	1016	11	0
1	G	1038	0	1015	1	0
1	H	1075	0	1060	5	0
2	A	28	0	26	2	0
2	B	28	0	26	1	0
2	C	42	0	39	5	0
2	D	14	0	13	0	0
2	E	28	0	26	2	0
2	F	28	0	26	2	0
2	G	28	0	26	4	0
2	H	28	0	26	1	0
3	A	100	0	0	2	0
3	B	92	0	0	3	0
3	C	89	0	0	0	0
3	D	72	0	0	1	0
3	E	78	0	0	0	0
3	F	86	0	0	2	0
3	G	80	0	0	1	0
3	H	85	0	0	0	0
All	All	9236	0	8356	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:502:NAG:C3	1:D:116[A]:ASN:HD22	1.82	0.92
2:G:502:NAG:O4	1:H:116[A]:ASN:ND2	2.05	0.89
2:C:502:NAG:O4	1:D:116[A]:ASN:ND2	2.16	0.77
1:F:130:LEU:HD21	1:F:132[B]:ILE:HD11	1.68	0.74
2:C:502:NAG:C3	1:D:116[A]:ASN:ND2	2.52	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/146 (86%)	123 (98%)	2 (2%)	0	100	100
1	B	127/146 (87%)	123 (97%)	4 (3%)	0	100	100
1	C	124/146 (85%)	122 (98%)	2 (2%)	0	100	100
1	D	124/146 (85%)	120 (97%)	4 (3%)	0	100	100
1	E	123/146 (84%)	119 (97%)	4 (3%)	0	100	100
1	F	126/146 (86%)	123 (98%)	3 (2%)	0	100	100
1	G	125/146 (86%)	123 (98%)	2 (2%)	0	100	100
1	H	130/146 (89%)	127 (98%)	3 (2%)	0	100	100
All	All	1004/1168 (86%)	980 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/138 (88%)	120 (99%)	1 (1%)	81	82
1	B	122/138 (88%)	122 (100%)	0	100	100
1	C	120/138 (87%)	120 (100%)	0	100	100
1	D	119/138 (86%)	119 (100%)	0	100	100
1	E	118/138 (86%)	117 (99%)	1 (1%)	81	82
1	F	122/138 (88%)	122 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	121/138 (88%)	118 (98%)	3 (2%)	47	41
1	H	125/138 (91%)	123 (98%)	2 (2%)	62	60
All	All	968/1104 (88%)	961 (99%)	7 (1%)	84	84

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	163	LYS
1	G	196	SER
1	H	196	SER
1	H	133	ARG
1	G	147	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	B	116	ASN
1	E	116	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	501	1	14,14,15	0.95	0	17,19,21	1.16	1 (5%)
2	NAG	G	502	1	14,14,15	1.17	2 (14%)	17,19,21	1.97	5 (29%)
2	NAG	F	501	1	14,14,15	0.70	0	17,19,21	1.19	1 (5%)
2	NAG	H	502	1	14,14,15	0.58	0	17,19,21	1.86	6 (35%)
2	NAG	H	501	1	14,14,15	1.34	2 (14%)	17,19,21	1.40	2 (11%)
2	NAG	C	502	1	14,14,15	1.11	2 (14%)	17,19,21	1.97	7 (41%)
2	NAG	D	501	1	14,14,15	0.80	0	17,19,21	1.38	3 (17%)
2	NAG	A	501	1	14,14,15	0.52	0	17,19,21	1.74	4 (23%)
2	NAG	E	502	1	14,14,15	0.94	1 (7%)	17,19,21	1.95	5 (29%)
2	NAG	C	501	1	14,14,15	1.02	1 (7%)	17,19,21	1.70	4 (23%)
2	NAG	B	502	1	14,14,15	1.04	2 (14%)	17,19,21	1.98	6 (35%)
2	NAG	G	501	1	14,14,15	1.11	1 (7%)	17,19,21	1.66	3 (17%)
2	NAG	E	501	1	14,14,15	0.94	1 (7%)	17,19,21	1.35	1 (5%)
2	NAG	A	502	1	14,14,15	1.06	2 (14%)	17,19,21	2.20	6 (35%)
2	NAG	C	503	1	14,14,15	0.34	0	17,19,21	2.08	5 (29%)
2	NAG	F	502	1	14,14,15	0.98	1 (7%)	17,19,21	1.89	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	501	1	-	0/6/23/26	0/1/1/1
2	NAG	G	502	1	-	0/6/23/26	0/1/1/1
2	NAG	F	501	1	-	0/6/23/26	0/1/1/1
2	NAG	H	502	1	-	0/6/23/26	0/1/1/1
2	NAG	H	501	1	-	0/6/23/26	0/1/1/1
2	NAG	C	502	1	-	0/6/23/26	0/1/1/1
2	NAG	D	501	1	-	0/6/23/26	0/1/1/1
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	E	502	1	-	0/6/23/26	0/1/1/1
2	NAG	C	501	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	502	1	-	0/6/23/26	0/1/1/1
2	NAG	G	501	1	-	1/6/23/26	0/1/1/1
2	NAG	E	501	1	-	0/6/23/26	0/1/1/1
2	NAG	A	502	1	-	0/6/23/26	0/1/1/1
2	NAG	C	503	1	-	2/6/23/26	0/1/1/1
2	NAG	F	502	1	-	0/6/23/26	0/1/1/1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	501	NAG	O5-C1	-3.28	1.38	1.43
2	G	502	NAG	O5-C1	-3.26	1.38	1.43
2	H	501	NAG	C2-N2	-2.85	1.41	1.46
2	C	501	NAG	O5-C1	-2.80	1.39	1.43
2	C	502	NAG	O5-C5	-2.55	1.38	1.43

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	503	NAG	C2-N2-C7	5.19	130.29	122.90
2	E	502	NAG	C4-C3-C2	4.62	117.79	111.02
2	G	501	NAG	O5-C1-C2	-4.45	104.26	111.29
2	E	501	NAG	C1-C2-N2	-4.44	102.91	110.49
2	A	502	NAG	C4-C3-C2	4.32	117.35	111.02

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	503	NAG	O5-C5-C6-O6
2	C	503	NAG	C4-C5-C6-O6
2	G	501	NAG	C4-C5-C6-O6

There are no ring outliers.

8 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	502	NAG	4	0
2	H	502	NAG	1	0
2	C	502	NAG	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NAG	1	0
2	E	502	NAG	2	0
2	B	502	NAG	1	0
2	A	502	NAG	1	0
2	F	502	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	124/146 (84%)	-0.40	0 100 100	17, 25, 47, 56	4 (3%)
1	B	128/146 (87%)	-0.62	0 100 100	18, 29, 46, 51	5 (3%)
1	C	126/146 (86%)	-0.66	1 (0%) 86 87	19, 26, 49, 73	6 (4%)
1	D	125/146 (85%)	-0.35	3 (2%) 59 62	19, 32, 61, 83	6 (4%)
1	E	125/146 (85%)	-0.61	1 (0%) 86 87	19, 31, 56, 78	11 (8%)
1	F	124/146 (84%)	-0.63	1 (0%) 86 87	18, 27, 52, 69	6 (4%)
1	G	124/146 (84%)	-0.52	1 (0%) 86 87	18, 28, 57, 86	8 (6%)
1	H	129/146 (88%)	-0.53	1 (0%) 86 87	19, 29, 52, 58	7 (5%)
All	All	1005/1168 (86%)	-0.54	8 (0%) 86 87	17, 28, 53, 86	53 (5%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	160	LEU	3.6
1	D	174	ASN	2.6
1	G	92	LEU	2.5
1	C	161	SER	2.4
1	H	216	PRO	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	502	14/15	0.69	0.23	30,36,38,40	14
2	NAG	H	502	14/15	0.69	0.30	31,37,41,45	14
2	NAG	B	502	14/15	0.74	0.27	29,37,42,45	14
2	NAG	A	502	14/15	0.77	0.32	27,32,37,43	14
2	NAG	C	502	14/15	0.77	0.20	21,32,37,40	14
2	NAG	C	503	14/15	0.80	0.18	23,36,42,44	14
2	NAG	G	502	14/15	0.81	0.29	26,29,36,45	14
2	NAG	F	502	14/15	0.82	0.22	25,31,38,39	14
2	NAG	E	501	14/15	0.93	0.07	28,33,39,41	0
2	NAG	G	501	14/15	0.94	0.09	27,36,53,57	0
2	NAG	F	501	14/15	0.94	0.07	24,32,38,43	0
2	NAG	D	501	14/15	0.94	0.10	30,39,45,45	0
2	NAG	C	501	14/15	0.96	0.06	27,32,36,40	0
2	NAG	A	501	14/15	0.97	0.07	23,28,40,41	0
2	NAG	H	501	14/15	0.97	0.07	20,22,27,28	0
2	NAG	B	501	14/15	0.97	0.07	22,25,28,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.