



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 3, 2023 – 06:44 AM EDT

PDB ID : 6PP2
Title : Structure of human endothelial nitric oxide synthase heme domain in complex with 4-((4-(2-Amino-4-methylquinolin-7-yl)-2-(aminomethyl)phenoxy)methyl)benzotrile
Authors : Chreifi, G.; Li, H.; Poulos, T.L.
Deposited on : 2019-07-05
Resolution : 2.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 14200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3237	2062	570	589	16	0	2	0
1	B	403	3232	2057	570	588	17	0	3	0
1	C	401	3209	2044	563	586	16	0	2	0
1	D	402	3221	2051	566	587	17	0	3	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	298	GLU	ASP	variant	UNP P29474
B	298	GLU	ASP	variant	UNP P29474
C	298	GLU	ASP	variant	UNP P29474
D	298	GLU	ASP	variant	UNP P29474

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		
3	C	1	Total	C	N	O	0	0
			17	9	5	3		
3	D	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 4-{{2-(aminomethyl)-4-(2-amino-4-methylquinolin-7-yl)phenoxy}methyl}benzotrile (three-letter code: OUM) (formula: C₂₅H₂₂N₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			30	25	4	1		
4	B	1	Total	C	N	O	0	0
			30	25	4	1		
4	C	1	Total	C	N	O	0	0
			30	25	4	1		
4	D	1	Total	C	N	O	0	0
			30	25	4	1		

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total 14	C 8	N 1	O 5	0	0
5	A	1	Total 14	C 8	N 1	O 5	0	0
5	B	1	Total 14	C 8	N 1	O 5	0	0
5	B	1	Total 14	C 8	N 1	O 5	0	0
5	B	1	Total 14	C 8	N 1	O 5	0	0
5	C	1	Total 14	C 8	N 1	O 5	0	0
5	C	1	Total 14	C 8	N 1	O 5	0	0
5	C	1	Total 14	C 8	N 1	O 5	0	0
5	D	1	Total 14	C 8	N 1	O 5	0	0
5	D	1	Total 14	C 8	N 1	O 5	0	0
5	D	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	Total 1	Zn 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Zn	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		
8	B	1	Total	Cl	0	0
			1	1		
8	C	1	Total	Cl	0	0
			1	1		
8	D	1	Total	Cl	0	0
			1	1		

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total 1	Gd 1	0	0
9	D	1	Total 1	Gd 1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	143	Total 143	O 143	0	0
10	B	239	Total 239	O 239	0	0
10	C	148	Total 148	O 148	0	0
10	D	237	Total 237	O 237	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.54Å 154.89Å 109.18Å 90.00° 90.80° 90.00°	Depositor
Resolution (Å)	89.23 – 2.02	Depositor
% Data completeness (in resolution range)	94.5 (89.23-2.02)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.02Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.199 , 0.246	Depositor
Wilson B-factor (Å ²)	27.9	Xtrriage
Anisotropy	0.649	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.105 for h,-k,-l	Xtrriage
Total number of atoms	14200	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 8 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BTB	C	504	-	13,13,13	2.91	1 (7%)	7,16,16	1.37	2 (28%)
5	BTB	B	506	-	13,13,13	0.64	0	7,16,16	1.21	1 (14%)
3	H4B	A	502	-	16,18,18	0.99	0	11,26,26	2.44	5 (45%)
7	GOL	A	507	-	5,5,5	0.35	0	5,5,5	0.26	0
3	H4B	B	502	-	16,18,18	0.98	0	11,26,26	2.63	6 (54%)
4	OUM	A	503	-	33,33,33	1.72	4 (12%)	45,46,46	2.17	12 (26%)
5	BTB	A	504	-	13,13,13	4.57	3 (23%)	7,16,16	2.23	2 (28%)
5	BTB	B	504	9	13,13,13	0.53	0	7,16,16	0.90	0
2	HEM	A	501	1	41,50,50	1.48	6 (14%)	45,82,82	1.82	9 (20%)
2	HEM	B	501	1	41,50,50	1.59	6 (14%)	45,82,82	1.79	9 (20%)
5	BTB	D	506	-	13,13,13	0.41	0	7,16,16	0.46	0
2	HEM	D	501	1	41,50,50	1.44	6 (14%)	45,82,82	1.96	12 (26%)
5	BTB	D	505	-	13,13,13	0.84	0	7,16,16	1.56	3 (42%)
5	BTB	B	505	-	13,13,13	0.57	0	7,16,16	0.65	0
3	H4B	D	502	-	16,18,18	0.91	0	11,26,26	2.54	5 (45%)
5	BTB	A	505	-	13,13,13	0.53	0	7,16,16	0.74	0
4	OUM	B	503	-	33,33,33	1.72	4 (12%)	45,46,46	1.88	14 (31%)
4	OUM	D	503	-	33,33,33	1.72	5 (15%)	45,46,46	2.04	12 (26%)
7	GOL	C	508	-	5,5,5	0.51	0	5,5,5	0.76	0
5	BTB	C	506	-	13,13,13	0.35	0	7,16,16	0.49	0
4	OUM	C	503	-	33,33,33	1.63	3 (9%)	45,46,46	1.89	9 (20%)
2	HEM	C	501	1	41,50,50	1.55	8 (19%)	45,82,82	1.84	9 (20%)
3	H4B	C	502	-	16,18,18	0.99	0	11,26,26	2.58	4 (36%)
5	BTB	C	505	-	13,13,13	1.02	2 (15%)	7,16,16	1.23	1 (14%)
5	BTB	D	504	9	13,13,13	0.37	0	7,16,16	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	C	504	-	-	7/21/21/21	-
5	BTB	B	506	-	-	9/21/21/21	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	502	-	-	0/8/17/17	0/2/2/2
7	GOL	A	507	-	-	4/4/4/4	-
3	H4B	B	502	-	-	0/8/17/17	0/2/2/2
4	OUM	A	503	-	-	1/13/13/13	0/4/4/4
5	BTB	A	504	-	-	14/21/21/21	-
5	BTB	B	504	9	-	2/21/21/21	-
2	HEM	A	501	1	-	3/12/54/54	-
2	HEM	B	501	1	-	2/12/54/54	-
5	BTB	D	506	-	-	14/21/21/21	-
2	HEM	D	501	1	-	3/12/54/54	-
5	BTB	D	505	-	-	13/21/21/21	-
5	BTB	B	505	-	-	5/21/21/21	-
3	H4B	D	502	-	-	3/8/17/17	0/2/2/2
5	BTB	A	505	-	-	4/21/21/21	-
4	OUM	B	503	-	-	2/13/13/13	0/4/4/4
4	OUM	D	503	-	-	2/13/13/13	0/4/4/4
7	GOL	C	508	-	-	2/4/4/4	-
5	BTB	C	506	-	-	3/21/21/21	-
4	OUM	C	503	-	-	5/13/13/13	0/4/4/4
2	HEM	C	501	1	-	5/12/54/54	-
3	H4B	C	502	-	-	0/8/17/17	0/2/2/2
5	BTB	C	505	-	-	5/21/21/21	-
5	BTB	D	504	9	-	1/21/21/21	-

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	504	BTB	C1-C2	-15.92	1.32	1.53
5	C	504	BTB	C5-N	-10.23	1.33	1.48
4	B	503	OUM	C34-C37	-7.70	1.27	1.44
4	A	503	OUM	C34-C37	-7.40	1.28	1.44
4	D	503	OUM	C34-C37	-7.35	1.28	1.44

The worst 5 of 115 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	OUM	C26-C21-C08	-7.78	108.00	120.86
4	C	503	OUM	O29-C24-C25	7.37	125.41	115.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	OUM	C22-C21-C08	6.70	132.96	121.36
2	B	501	HEM	CBA-CAA-C2A	-6.32	101.84	112.62
3	C	502	H4B	C8A-C4A-C4	6.03	119.93	114.57

There are no chirality outliers.

5 of 109 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C2A-CAA-CBA-CGA
2	C	501	HEM	C2D-C3D-CAD-CBD
2	C	501	HEM	C4D-C3D-CAD-CBD
4	B	503	OUM	C24-C25-C27-N28
5	A	504	BTB	O1-C1-C2-C4


There are no ring outliers.


1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	505	BTB	0	1


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.