



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 03:25 pm BST

PDB ID : 1TRP
Title : X-RAY CRYSTALLOGRAPHIC AND CALORIMERIC STUDIES OF THE EFFECTS OF THE MUTATION TRP 59 TYR IN RIBONUCLEASE T1
Authors : Schluckebier, G.; Saenger, W.
Deposited on : 1993-07-06
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	104	 67% 28% . .
1	B	104	 66% 28% 6%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 1704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE T1 ISOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	104	778	479	127	168	4	0	0	0
1	B	104	783	481	127	171	4	0	1	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	LYS	GLN	CONFLICT	UNP P00651
A	45	TRP	TYR	CONFLICT	UNP P00651
A	59	TYR	TRP	CONFLICT	UNP P00651
B	225	LYS	GLN	CONFLICT	UNP P00651
B	245	TRP	TYR	CONFLICT	UNP P00651
B	259	TYR	TRP	CONFLICT	UNP P00651

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is GUANOSINE-2'-MONOPHOSPHATE (three-letter code: 2GP) (formula: C₁₀H₁₄N₅O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	24	10	5	8	1	0	0
3	B	1	24	10	5	8	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	55	55	55	0	0
4	B	38	38	38	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBONUCLEASE T1 ISOZYME



- Molecule 1: RIBONUCLEASE T1 ISOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.15Å 48.15Å 40.15Å 90.00° 90.35° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40 14.83 – 2.38	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.40) 83.4 (14.83-2.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.96 (at 2.39Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.160 , (Not available) 0.250 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 61.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l 0.035 for -k,-h,-l 0.186 for -h,-k,l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	1704	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2GP, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/799	0.94	0/1089
1	B	0.62	0/810	0.96	0/1104
All	All	0.62	0/1609	0.95	0/2193

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	778	0	674	31	1
1	B	783	0	673	29	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	24	0	12	2	0
3	B	24	0	12	2	0
4	A	55	0	0	0	0
4	B	38	0	0	3	0
All	All	1704	0	1371	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ALA:HB2	1:A:101:VAL:HG13	1.28	1.16
1:B:252:VAL:HG21	1:B:285:GLN:HB3	1.53	0.87
1:A:50:PHE:HD2	1:A:56:TYR:CZ	2.05	0.73
1:B:252:VAL:HG21	1:B:285:GLN:CB	2.21	0.71
1:A:95:ALA:CB	1:A:101:VAL:HG13	2.15	0.70
1:A:32:THR:HG21	1:A:37:SER:HB3	1.74	0.69
1:A:81:ASN:HD21	1:A:85:GLN:NE2	1.91	0.69
1:B:254:SER:HB3	1:B:255:PRO:HA	1.74	0.69
1:A:95:ALA:HB2	1:A:101:VAL:CG1	2.16	0.68
1:B:279:VAL:HG21	1:B:290:ILE:HD13	1.80	0.63
1:B:235:SER:HB2	4:B:492:HOH:O	1.98	0.63
1:A:81:ASN:HD21	1:A:85:GLN:HE21	1.43	0.62
1:B:252:VAL:HG22	1:B:281:ASN:HD21	1.64	0.62
3:B:305:2GP:H3'	3:B:305:2GP:N3	2.16	0.61
1:A:83:ASN:ND2	1:B:245:TRP:HE1	1.98	0.61
1:B:279:VAL:CG2	1:B:290:ILE:HD13	2.30	0.60
1:B:252:VAL:HG22	1:B:281:ASN:ND2	2.16	0.60
1:B:254:SER:CB	1:B:255:PRO:HA	2.33	0.58
1:A:54:SER:OG	1:A:55:PRO:HA	2.03	0.57
1:A:52:VAL:HG21	1:A:85:GLN:CB	2.34	0.57
1:A:83:ASN:HD21	1:B:245:TRP:HZ2	1.51	0.57
1:B:303:CYS:O	1:B:304:THR:HB	2.05	0.57
1:A:16:VAL:HG11	1:A:86:LEU:HD11	1.87	0.56
1:A:80:PHE:HA	1:A:85:GLN:O	2.06	0.56
1:B:204:TYR:CD2	1:B:216:VAL:HG21	2.41	0.56
1:A:81:ASN:ND2	1:A:85:GLN:HE21	2.05	0.55
1:B:205:THR:HG22	4:B:454:HOH:O	2.07	0.54
3:A:105:2GP:N3	3:A:105:2GP:H3'	2.23	0.53
1:B:281:ASN:HD21	1:B:285:GLN:NE2	2.07	0.52
1:B:274:GLY:HA3	4:B:444:HOH:O	2.09	0.51
1:B:296:SER:O	1:B:299:ASN:ND2	2.43	0.51
1:A:58:GLU:OE1	3:A:105:2GP:O2P	2.29	0.51
1:A:32:THR:CG2	1:A:37:SER:HB3	2.39	0.49
1:A:52:VAL:HG21	1:A:85:GLN:HB2	1.93	0.49
1:A:92:HIS:O	1:A:95:ALA:HB3	2.13	0.48
1:A:96:SER:O	1:A:99:ASN:HB2	2.15	0.47
1:B:244:ASN:N	3:B:305:2GP:O6	2.49	0.46
1:A:83:ASN:HD22	1:B:245:TRP:HE1	1.62	0.46
1:A:90:ILE:HD12	1:A:100:PHE:CG	2.51	0.45
1:B:279:VAL:HG12	1:B:287:ALA:HB3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ASN:HD21	1:A:48:PHE:H	1.65	0.45
1:B:281:ASN:HD21	1:B:285:GLN:HE21	1.65	0.43
1:B:226:LEU:HD13	1:B:233:VAL:HG22	2.01	0.43
1:B:292:HIS:HE1	1:B:298:ASN:O	2.02	0.43
1:A:59:TYR:CG	1:A:60:PRO:HD2	2.53	0.42
1:A:52:VAL:HG21	1:A:85:GLN:HB3	2.01	0.42
1:A:76:ASP:OD1	1:A:91:THR:OG1	2.35	0.42
1:B:254:SER:HB3	1:B:255:PRO:CA	2.46	0.42
1:A:44:ASN:HA	1:A:44:ASN:HD22	1.62	0.42
1:A:90:ILE:HD12	1:A:100:PHE:CD1	2.55	0.41
1:B:205:THR:HG23	1:B:304:THR:HA	2.02	0.41
1:A:59:TYR:CD1	1:A:60:PRO:HD2	2.55	0.41
1:B:286:LEU:HD22	1:B:288:GLY:H	1.86	0.41
1:A:50:PHE:HD2	1:A:56:TYR:CE2	2.39	0.41
1:B:206:CYS:HB3	1:B:291:THR:HB	2.02	0.40
1:A:54:SER:CB	1:A:55:PRO:HA	2.51	0.40
1:A:83:ASN:ND2	1:B:245:TRP:NE1	2.65	0.40
1:B:285:GLN:HE21	1:B:285:GLN:HB2	1.64	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:GLY:O	1:B:263:SER:OG[2_656]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/104 (98%)	95 (93%)	6 (6%)	1 (1%)	15	23
1	B	103/104 (99%)	100 (97%)	2 (2%)	1 (1%)	15	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	205/208 (99%)	195 (95%)	8 (4%)	2 (1%)	15	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	B	237	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/85 (99%)	73 (87%)	11 (13%)	4	4
1	B	86/85 (101%)	73 (85%)	13 (15%)	3	3
All	All	170/170 (100%)	146 (86%)	24 (14%)	3	4

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	SER
1	A	13	SER
1	A	25	LYS
1	A	33	VAL
1	A	35	SER
1	A	41	LYS
1	A	44	ASN
1	A	52	VAL
1	A	54	SER
1	A	83	ASN
1	A	86	LEU
1	B	205	THR
1	B	208	SER
1	B	217	SER
1	B	231[A]	GLU
1	B	231[B]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	233	VAL
1	B	235	SER
1	B	251	SER
1	B	252	VAL
1	B	278	VAL
1	B	284	ASN
1	B	286	LEU
1	B	304	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	44	ASN
1	A	83	ASN
1	A	84	ASN
1	A	85	GLN
1	B	220	GLN
1	B	243	ASN
1	B	283	ASN
1	B	284	ASN
1	B	285	GLN
1	B	299	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2GP	B	305	-	21,26,26	1.09	2 (9%)	26,40,40	2.31	4 (15%)
3	2GP	A	105	-	21,26,26	1.15	1 (4%)	26,40,40	2.25	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2GP	B	305	-	-	4/7/27/27	0/3/3/3
3	2GP	A	105	-	-	3/7/27/27	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	105	2GP	C6-N1	4.21	1.40	1.33
3	B	305	2GP	C6-N1	3.86	1.39	1.33
3	B	305	2GP	C8-N7	-2.14	1.30	1.34

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	105	2GP	C5-C6-N1	-8.61	111.65	123.43
3	B	305	2GP	C5-C6-N1	-8.47	111.85	123.43
3	B	305	2GP	C6-N1-C2	5.67	124.95	115.93
3	A	105	2GP	C6-N1-C2	5.54	124.74	115.93
3	B	305	2GP	C2-N3-C4	-3.28	111.61	115.36
3	A	105	2GP	C2-N3-C4	-3.14	111.78	115.36
3	A	105	2GP	N3-C2-N1	-2.42	123.99	127.22
3	B	305	2GP	N3-C2-N1	-2.41	124.01	127.22

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	305	2GP	C2'-O2'-P-O1P
3	B	305	2GP	C3'-C4'-C5'-O5'
3	B	305	2GP	O4'-C4'-C5'-O5'
3	A	105	2GP	C3'-C4'-C5'-O5'
3	A	105	2GP	O4'-C4'-C5'-O5'
3	A	105	2GP	C2'-O2'-P-O3P
3	B	305	2GP	C2'-O2'-P-O3P

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	305	2GP	2	0
3	A	105	2GP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands


Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2GP A 105:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.