



wwPDB EM Validation Summary Report ⓘ

Jun 22, 2023 – 11:03 AM JST

PDB ID : 7VYN
EMDB ID : EMD-32210
Title : Matrix arm of active state CI from Q1-NADH dataset
Authors : Gu, J.; Yang, M.
Deposited on : 2021-11-14
Resolution : 2.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev50
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|--------------------------|--------------------------|
| Clashscore | 158937 | 4297 |
| Ramachandran outliers | 154571 | 4023 |
| Sidechain outliers | 154315 | 3826 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 433 | |
| 2 | B | 176 | |
| 3 | C | 156 | |
| 4 | E | 115 | |
| 5 | F | 86 | |
| 6 | G | 88 | |
| 7 | H | 112 | |
| 8 | I | 112 | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 9 | J | 342 |  |
| 10 | K | 43 |  |
| 11 | L | 125 |  |
| 12 | M | 690 |  |
| 13 | N | 144 |  |
| 14 | O | 217 |  |
| 15 | P | 208 |  |
| 16 | Q | 386 |  |
| 17 | T | 96 |  |
| 18 | W | 29 |  |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 19 | SF4 | A | 501 | - | - | X | - |

2 Entry composition [i](#)

There are 32 unique types of molecules in this entry. The entry contains 29298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 1 | A | 433 | 3330 | 2103 | 593 | 614 | 20 | 0 | 0 |

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 2 | B | 176 | 1412 | 887 | 243 | 269 | 13 | 0 | 0 |

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 3 | C | 156 | 1248 | 794 | 227 | 213 | 14 | 0 | 0 |

- Molecule 4 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 4 | E | 115 | 971 | 619 | 179 | 168 | 5 | 0 | 0 |

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 5 | F | 86 | 687 | 432 | 129 | 124 | 2 | 0 | 0 |

- Molecule 6 is a protein called Acyl carrier protein, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 6 | G | 88 | 693 | 447 | 102 | 139 | 5 | 0 | 0 |

- Molecule 7 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 7 | H | 112 | 910 | 588 | 154 | 165 | 3 | 0 | 0 |

- Molecule 8 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 8 | I | 97 | 780 | 491 | 147 | 139 | 3 | 0 | 0 |

- Molecule 9 is a protein called NADH dehydrogenase ubiquinone 1 alpha subcomplex subunit 9, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 9 | J | 342 | 2751 | 1783 | 481 | 478 | 9 | 0 | 0 |

- Molecule 10 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 10 | K | 43 | 366 | 228 | 68 | 69 | 1 | 0 | 0 |

- Molecule 11 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 11 | L | 125 | 1016 | 642 | 181 | 190 | 3 | 0 | 0 |

- Molecule 12 is a protein called NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|------|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 12 | M | 690 | 5296 | 3320 | 923 | 1014 | 39 | 0 | 0 |

- Molecule 13 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 13 | N | 144 | 1204 | 770 | 218 | 212 | 4 | 0 | 0 |

- Molecule 14 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 14 | O | 217 | 1671 | 1065 | 281 | 315 | 10 | 0 | 0 |

- Molecule 15 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 3, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 15 | P | 208 | 1738 | 1124 | 298 | 314 | 2 | 0 | 0 |

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 16 | Q | 386 | 3096 | 1976 | 534 | 563 | 23 | 0 | 0 |


- Molecule 17 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 17 | T | 96 | 741 | 452 | 140 | 146 | 3 | 0 | 0 |

- Molecule 18 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13.


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 18 | W | 29 | 218 | 138 | 40 | 39 | 1 | 0 | 0 |

- Molecule 19 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | AltConf |
|-----|-------|----------|-------|----|---|---------|
| | | | Total | Fe | S | |
| 19 | A | 1 | 8 | 4 | 4 | 0 |
| 19 | B | 1 | 8 | 4 | 4 | 0 |
| 19 | B | 1 | 8 | 4 | 4 | 0 |
| 19 | C | 1 | 8 | 4 | 4 | 0 |
| 19 | M | 1 | 8 | 4 | 4 | 0 |
| 19 | M | 1 | 8 | 4 | 4 | 0 |

- Molecule 20 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|-----|-------|----------|-------|----|---|---|---|---------|
| | | | Total | C | N | O | P | |
| 20 | A | 1 | 31 | 17 | 4 | 9 | 1 | 0 |

- Molecule 21 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: $C_{21}H_{29}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|-----|-------|----------|-------|----|---|----|---|---------|
| | | | Total | C | N | O | P | |
| 21 | A | 1 | 44 | 21 | 7 | 14 | 2 | 0 |

- Molecule 22 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (three-letter code: PEE) (formula: $C_{41}H_{78}NO_8P$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|-----|-------|----------|-------|----|---|---|---|---------|
| | | | Total | C | N | O | P | |
| 22 | C | 1 | 47 | 37 | 1 | 8 | 1 | 0 |

- Molecule 23 is (9R,11S)-9-({[(1S)-1-HYDROXYHEXADECYL]OXY}METHYL)-2,2-DIMETHYL-5,7,10-TRIOXA-2LAMBDA 5 -AZA-6LAMBDA 5 -PHOSPHAOCTACOSANE-6,6,11-TRIOXANE (three-letter code: PLX) (formula: C₄₂H₈₉NO₈P) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf |
|-----|-------|----------|-------|----|---|---|---|---------|
| | | | Total | C | N | O | P | |
| 23 | C | 1 | 52 | 42 | 1 | 8 | 1 | 0 |

- Molecule 24 is S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonoxy)butanoyl]-beta-alanyl}amino)ethyl] dodecanethioate (three-letter code: 8Q1) (formula: $C_{23}H_{45}N_2O_8PS$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | | AltConf | |
|-----|-------|----------|-------|----|---|---|---|---------|---|
| | | | Total | C | N | O | P | | S |
| 24 | E | 1 | 35 | 23 | 2 | 8 | 1 | 1 | 0 |

- Molecule 25 is CARDIOLIPIN (three-letter code: CDL) (formula: $C_{81}H_{156}O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|----|----|---|---------|
| | | | Total | C | O | P | |
| 25 | I | 1 | 51 | 32 | 17 | 2 | 0 |

- Molecule 26 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | | AltConf | |
|-----|-------|----------|-------|----|---|----|---------|---|
| | | | Total | C | N | O | | P |
| 26 | J | 1 | 48 | 21 | 7 | 17 | 3 | 0 |

- Molecule 27 is Coenzyme Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-isomer (three-letter code: UQ) (formula: $C_{59}H_{90}O_4$) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | AltConf |
|-----|-------|----------|-------|----|---|---------|
| 27 | J | 1 | Total | C | O | 0 |
| | | | 33 | 29 | 4 | |

- Molecule 28 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | AltConf |
|-----|-------|----------|-------|----|---|---------|
| 28 | M | 1 | Total | Fe | S | 0 |
| | | | 4 | 2 | 2 | |
| 28 | O | 1 | Total | Fe | S | 0 |
| | | | 4 | 2 | 2 | |

- Molecule 29 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms | AltConf |
|-----|-------|----------|-----------------|---------|
| 29 | M | 1 | Total Mg 1 1 | 0 |

- Molecule 30 is UBIQUINONE-1 (three-letter code: UQ1) (formula: C₁₄H₁₈O₄) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | AltConf |
|-----|-------|----------|----------------------|---------|
| 30 | Q | 1 | Total C O 18 14 4 | 0 |
| 30 | Q | 1 | Total C O 18 14 4 | 0 |

- Molecule 31 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms | AltConf |
|-----|-------|----------|-----------------|---------|
| 31 | T | 1 | Total Zn 1 1 | 0 |

- Molecule 32 is water.

| Mol | Chain | Residues | Atoms | AltConf |
|-----|-------|----------|------------------|---------|
| 32 | A | 50 | Total O 50 50 | 0 |

Continued on next page...


Continued from previous page...

| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|--------------|----------|---------|
| 32 | B | 76 | Total 76 | O 76 | 0 |
| 32 | C | 54 | Total 54 | O 54 | 0 |
| 32 | E | 3 | Total 3 | O 3 | 0 |
| 32 | F | 1 | Total 1 | O 1 | 0 |
| 32 | H | 6 | Total 6 | O 6 | 0 |
| 32 | I | 14 | Total 14 | O 14 | 0 |
| 32 | J | 7 | Total 7 | O 7 | 0 |
| 32 | K | 6 | Total 6 | O 6 | 0 |
| 32 | L | 24 | Total 24 | O 24 | 0 |
| 32 | M | 196 | Total 196 | O 196 | 0 |
| 32 | N | 8 | Total 8 | O 8 | 0 |
| 32 | O | 14 | Total 14 | O 14 | 0 |
| 32 | P | 74 | Total 74 | O 74 | 0 |
| 32 | Q | 191 | Total 191 | O 191 | 0 |
| 32 | T | 9 | Total 9 | O 9 | 0 |
| 32 | W | 2 | Total 2 | O 2 | 0 |


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial




- Molecule 2: NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial




- Molecule 3: NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial




- Molecule 4: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6




- Molecule 5: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2




- Molecule 6: Acyl carrier protein, mitochondrial




- Molecule 7: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5




- Molecule 8: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7



- Molecule 9: NADH dehydrogenase ubiquinone 1 alpha subcomplex subunit 9, mitochondrial




- Molecule 10: NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial







- Molecule 11: NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial





- Molecule 12: NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial



- Molecule 13: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12



- Molecule 14: NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial




- Molecule 15: NADH dehydrogenase [ubiquinone] iron-sulfur protein 3, mitochondrial

Chain P:  88% 12%




- Molecule 16: NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial

Chain Q:  86% 14%




- Molecule 17: NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial

Chain T:  7% 91% 9%



- Molecule 18: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13

Chain W:  24% 93% 7%



4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | |
| Number of particles used | 302353 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 50 | Depositor |
| Minimum defocus (nm) | 1300 | Depositor |
| Maximum defocus (nm) | 1800 | Depositor |
| Magnification | Not provided | |
| Image detector | GATAN K3 (6k x 4k) | Depositor |
| Maximum map value | 0.127 | Depositor |
| Minimum map value | -0.065 | Depositor |
| Average map value | 0.000 | Depositor |
| Map value standard deviation | 0.003 | Depositor |
| Recommended contour level | 0.0162 | Depositor |
| Map size (Å) | 263.496, 263.496, 263.496 | wwPDB |
| Map dimensions | 480, 480, 480 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 0.54895, 0.54895, 0.54895 | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEE, SF4, MG, NAI, 8Q1, 2MR, ZN, NDP, UQ, CDL, FES, PLX, FMN, UQ1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.26 | 0/3406 | 0.50 | 0/4603 |
| 2 | B | 0.26 | 0/1443 | 0.51 | 0/1952 |
| 3 | C | 0.27 | 0/1279 | 0.54 | 0/1730 |
| 4 | E | 0.24 | 0/995 | 0.50 | 0/1340 |
| 5 | F | 0.25 | 0/698 | 0.56 | 0/940 |
| 6 | G | 0.24 | 0/705 | 0.41 | 0/956 |
| 7 | H | 0.25 | 0/929 | 0.46 | 0/1258 |
| 8 | I | 0.26 | 0/798 | 0.55 | 0/1079 |
| 9 | J | 0.25 | 0/2828 | 0.49 | 0/3834 |
| 10 | K | 0.25 | 0/377 | 0.51 | 0/509 |
| 11 | L | 0.24 | 0/1039 | 0.51 | 0/1403 |
| 12 | M | 0.25 | 0/5384 | 0.51 | 1/7295 (0.0%) |
| 13 | N | 0.25 | 0/1245 | 0.51 | 0/1694 |
| 14 | O | 0.26 | 0/1711 | 0.50 | 0/2328 |
| 15 | P | 0.27 | 0/1789 | 0.51 | 0/2436 |
| 16 | Q | 0.27 | 0/3157 | 0.51 | 0/4268 |
| 17 | T | 0.25 | 0/755 | 0.53 | 0/1018 |
| 18 | W | 0.26 | 0/224 | 0.51 | 0/302 |
| All | All | 0.26 | 0/28762 | 0.51 | 1/38945 (0.0%) |

There are no bond length outliers.

All (1) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|------|-------------|----------|
| 12 | M | 563 | ASP | CB-CG-OD1 | 5.11 | 122.90 | 118.30 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 3330 | 0 | 3292 | 44 | 0 |
| 2 | B | 1412 | 0 | 1363 | 13 | 0 |
| 3 | C | 1248 | 0 | 1254 | 17 | 0 |
| 4 | E | 971 | 0 | 975 | 18 | 0 |
| 5 | F | 687 | 0 | 700 | 14 | 0 |
| 6 | G | 693 | 0 | 671 | 12 | 0 |
| 7 | H | 910 | 0 | 950 | 7 | 0 |
| 8 | I | 780 | 0 | 808 | 12 | 0 |
| 9 | J | 2751 | 0 | 2773 | 29 | 0 |
| 10 | K | 366 | 0 | 338 | 5 | 0 |
| 11 | L | 1016 | 0 | 1016 | 15 | 0 |
| 12 | M | 5296 | 0 | 5326 | 62 | 0 |
| 13 | N | 1204 | 0 | 1162 | 18 | 0 |
| 14 | O | 1671 | 0 | 1673 | 22 | 0 |
| 15 | P | 1738 | 0 | 1693 | 16 | 0 |
| 16 | Q | 3096 | 0 | 3063 | 31 | 0 |
| 17 | T | 741 | 0 | 702 | 5 | 0 |
| 18 | W | 218 | 0 | 219 | 2 | 0 |
| 19 | A | 8 | 0 | 0 | 2 | 0 |
| 19 | B | 16 | 0 | 0 | 0 | 0 |
| 19 | C | 8 | 0 | 0 | 1 | 0 |
| 19 | M | 16 | 0 | 0 | 0 | 0 |
| 20 | A | 31 | 0 | 19 | 4 | 0 |
| 21 | A | 44 | 0 | 27 | 5 | 0 |
| 22 | C | 47 | 0 | 71 | 4 | 0 |
| 23 | C | 52 | 0 | 88 | 3 | 0 |
| 24 | E | 35 | 0 | 0 | 4 | 0 |
| 25 | I | 51 | 0 | 46 | 1 | 0 |
| 26 | J | 48 | 0 | 25 | 2 | 0 |
| 27 | J | 33 | 0 | 39 | 2 | 0 |
| 28 | M | 4 | 0 | 0 | 0 | 0 |
| 28 | O | 4 | 0 | 0 | 0 | 0 |
| 29 | M | 1 | 0 | 0 | 0 | 0 |
| 30 | Q | 36 | 0 | 36 | 2 | 0 |
| 31 | T | 1 | 0 | 0 | 0 | 0 |
| 32 | A | 50 | 0 | 0 | 1 | 0 |
| 32 | B | 76 | 0 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 32 | C | 54 | 0 | 0 | 0 | 0 |
| 32 | E | 3 | 0 | 0 | 0 | 0 |
| 32 | F | 1 | 0 | 0 | 0 | 0 |
| 32 | H | 6 | 0 | 0 | 0 | 0 |
| 32 | I | 14 | 0 | 0 | 0 | 0 |
| 32 | J | 7 | 0 | 0 | 0 | 0 |
| 32 | K | 6 | 0 | 0 | 0 | 0 |
| 32 | L | 24 | 0 | 0 | 0 | 0 |
| 32 | M | 196 | 0 | 0 | 1 | 0 |
| 32 | N | 8 | 0 | 0 | 0 | 0 |
| 32 | O | 14 | 0 | 0 | 0 | 0 |
| 32 | P | 74 | 0 | 0 | 0 | 0 |
| 32 | Q | 191 | 0 | 0 | 5 | 0 |
| 32 | T | 9 | 0 | 0 | 0 | 0 |
| 32 | W | 2 | 0 | 0 | 0 | 0 |
| All | All | 29298 | 0 | 28329 | 296 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-------------------|-------------------|--------------------------|-------------------|
| 24:E:201:8Q1:O1 | 6:G:112:SER:CB | 1.86 | 1.23 |
| 26:J:401:NDP:O4D | 26:J:401:NDP:C4D | 1.69 | 1.16 |
| 21:A:503:NAI:C1B | 21:A:503:NAI:O4B | 1.63 | 1.16 |
| 14:O:164:THR:HG22 | 14:O:165:PRO:HD2 | 1.29 | 1.09 |
| 1:A:121:GLU:HB2 | 21:A:503:NAI:H42N | 1.59 | 0.83 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 1 | A | 431/433 (100%) | 420 (97%) | 11 (3%) | 0 | 100 | 100 |
| 2 | B | 174/176 (99%) | 171 (98%) | 3 (2%) | 0 | 100 | 100 |
| 3 | C | 154/156 (99%) | 150 (97%) | 4 (3%) | 0 | 100 | 100 |
| 4 | E | 113/115 (98%) | 107 (95%) | 6 (5%) | 0 | 100 | 100 |
| 5 | F | 84/86 (98%) | 81 (96%) | 3 (4%) | 0 | 100 | 100 |
| 6 | G | 86/88 (98%) | 83 (96%) | 3 (4%) | 0 | 100 | 100 |
| 7 | H | 110/112 (98%) | 102 (93%) | 7 (6%) | 1 (1%) | 17 | 25 |
| 8 | I | 93/112 (83%) | 82 (88%) | 11 (12%) | 0 | 100 | 100 |
| 9 | J | 340/342 (99%) | 329 (97%) | 10 (3%) | 1 (0%) | 41 | 55 |
| 10 | K | 41/43 (95%) | 40 (98%) | 1 (2%) | 0 | 100 | 100 |
| 11 | L | 123/125 (98%) | 121 (98%) | 2 (2%) | 0 | 100 | 100 |
| 12 | M | 688/690 (100%) | 665 (97%) | 23 (3%) | 0 | 100 | 100 |
| 13 | N | 142/144 (99%) | 138 (97%) | 4 (3%) | 0 | 100 | 100 |
| 14 | O | 215/217 (99%) | 206 (96%) | 9 (4%) | 0 | 100 | 100 |
| 15 | P | 206/208 (99%) | 197 (96%) | 9 (4%) | 0 | 100 | 100 |
| 16 | Q | 383/386 (99%) | 372 (97%) | 11 (3%) | 0 | 100 | 100 |
| 17 | T | 94/96 (98%) | 93 (99%) | 1 (1%) | 0 | 100 | 100 |
| 18 | W | 27/29 (93%) | 24 (89%) | 3 (11%) | 0 | 100 | 100 |
| All | All | 3504/3558 (98%) | 3381 (96%) | 121 (4%) | 2 (0%) | 54 | 68 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 7 | H | 77 | ILE |
| 9 | J | 38 | HIS |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|-----|
| 1 | A | 346/346 (100%) | 335 (97%) | 11 (3%) | 39 | 59 |
| 2 | B | 151/151 (100%) | 149 (99%) | 2 (1%) | 69 | 84 |
| 3 | C | 132/132 (100%) | 127 (96%) | 5 (4%) | 33 | 51 |
| 4 | E | 107/107 (100%) | 103 (96%) | 4 (4%) | 34 | 53 |
| 5 | F | 75/76 (99%) | 73 (97%) | 2 (3%) | 44 | 65 |
| 6 | G | 76/81 (94%) | 73 (96%) | 3 (4%) | 32 | 50 |
| 7 | H | 99/99 (100%) | 96 (97%) | 3 (3%) | 41 | 61 |
| 8 | I | 87/97 (90%) | 81 (93%) | 6 (7%) | 15 | 25 |
| 9 | J | 296/296 (100%) | 291 (98%) | 5 (2%) | 60 | 78 |
| 10 | K | 42/42 (100%) | 40 (95%) | 2 (5%) | 25 | 41 |
| 11 | L | 113/113 (100%) | 110 (97%) | 3 (3%) | 44 | 65 |
| 12 | M | 580/580 (100%) | 561 (97%) | 19 (3%) | 38 | 57 |
| 13 | N | 130/130 (100%) | 123 (95%) | 7 (5%) | 22 | 36 |
| 14 | O | 183/183 (100%) | 177 (97%) | 6 (3%) | 38 | 57 |
| 15 | P | 190/190 (100%) | 189 (100%) | 1 (0%) | 88 | 95 |
| 16 | Q | 332/332 (100%) | 322 (97%) | 10 (3%) | 41 | 61 |
| 17 | T | 79/79 (100%) | 77 (98%) | 2 (2%) | 47 | 67 |
| 18 | W | 23/24 (96%) | 23 (100%) | 0 | 100 | 100 |
| All | All | 3041/3058 (99%) | 2950 (97%) | 91 (3%) | 44 | 61 |

5 of 91 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 12 | M | 467 | LYS |
| 13 | N | 132 | LYS |
| 12 | M | 581 | ASP |
| 13 | N | 9 | ARG |
| 14 | O | 164 | THR |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 204 | ASN |
| 4 | E | 70 | ASN |
| 11 | L | 109 | ASN |
| 14 | O | 153 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 16 | 2MR | Q | 118 | 16 | 10,12,13 | 2.01 | 1 (10%) | 5,13,15 | 5.92 | 3 (60%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|------------|-------|
| 16 | 2MR | Q | 118 | 16 | - | 2/10/13/15 | - |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 16 | Q | 118 | 2MR | CZ-NE | 5.73 | 1.46 | 1.34 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^{\circ}$) | Ideal($^{\circ}$) |
|-----|-------|-----|------|------------|-------|------------------------|---------------------|
| 16 | Q | 118 | 2MR | NE-CZ-NH2 | 12.02 | 130.50 | 119.48 |
| 16 | Q | 118 | 2MR | CD-NE-CZ | 3.77 | 130.47 | 123.41 |
| 16 | Q | 118 | 2MR | CQ2-NH2-CZ | 3.57 | 131.76 | 123.86 |

There are no chirality outliers.

All (2) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 16 | Q | 118 | 2MR | NE-CD-CG-CB |
| 16 | Q | 118 | 2MR | CA-CB-CG-CD |

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 2 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 19 | SF4 | B | 302 | 2 | 0,12,12 | - | - | - | | |
| 30 | UQ1 | Q | 501 | - | 18,18,18 | 2.38 | 5 (27%) | 22,25,25 | 1.95 | 6 (27%) |
| 21 | NAI | A | 503 | - | 42,48,48 | 4.93 | 18 (42%) | 47,73,73 | 1.35 | 7 (14%) |
| 25 | CDL | I | 201 | - | 50,50,99 | 1.40 | 8 (16%) | 56,62,111 | 1.13 | 4 (7%) |
| 19 | SF4 | M | 801 | 12 | 0,12,12 | - | - | - | | |
| 19 | SF4 | M | 802 | 12 | 0,12,12 | - | - | - | | |
| 23 | PLX | C | 303 | - | 51,51,51 | 1.12 | 3 (5%) | 55,59,59 | 0.59 | 1 (1%) |
| 30 | UQ1 | Q | 502 | - | 18,18,18 | 2.33 | 6 (33%) | 22,25,25 | 1.59 | 4 (18%) |
| 20 | FMN | A | 502 | - | 33,33,33 | 1.10 | 2 (6%) | 48,50,50 | 1.28 | 9 (18%) |
| 26 | NDP | J | 401 | - | 45,52,52 | 4.56 | 20 (44%) | 53,80,80 | 1.92 | 7 (13%) |
| 19 | SF4 | A | 501 | 1 | 0,12,12 | - | - | - | | |
| 24 | 8Q1 | E | 201 | - | 31,34,34 | 1.70 | 6 (19%) | 40,43,43 | 1.61 | 6 (15%) |
| 27 | UQ | J | 402 | - | 33,33,63 | 3.45 | 10 (30%) | 40,43,79 | 2.72 | 14 (35%) |
| 19 | SF4 | C | 301 | 3 | 0,12,12 | - | - | - | | |
| 19 | SF4 | B | 301 | 2 | 0,12,12 | - | - | - | | |
| 22 | PEE | C | 302 | - | 46,46,50 | 1.21 | 6 (13%) | 49,51,55 | 0.95 | 2 (4%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 28 | FES | M | 803 | 12 | 0,4,4 | - | - | - | | |
| 28 | FES | O | 301 | 14 | 0,4,4 | - | - | - | | |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|--------------|---------|
| 30 | UQ1 | Q | 501 | - | - | 4/9/33/33 | 0/1/1/1 |
| 19 | SF4 | B | 302 | 2 | - | - | 0/6/5/5 |
| 21 | NAI | A | 503 | - | - | 6/25/72/72 | 0/5/5/5 |
| 25 | CDL | I | 201 | - | - | 30/61/61/110 | - |
| 23 | PLX | C | 303 | - | - | 25/55/55/55 | - |
| 19 | SF4 | M | 801 | 12 | - | - | 0/6/5/5 |
| 19 | SF4 | M | 802 | 12 | - | - | 0/6/5/5 |
| 30 | UQ1 | Q | 502 | - | - | 2/9/33/33 | 0/1/1/1 |
| 20 | FMN | A | 502 | - | - | 9/18/18/18 | 0/3/3/3 |
| 26 | NDP | J | 401 | - | - | 10/30/77/77 | 0/4/5/5 |
| 27 | UQ | J | 402 | - | - | 13/27/51/87 | 0/1/1/1 |
| 19 | SF4 | A | 501 | 1 | - | - | 0/6/5/5 |
| 22 | PEE | C | 302 | - | - | 25/50/50/54 | - |
| 19 | SF4 | B | 301 | 2 | - | - | 0/6/5/5 |
| 24 | 8Q1 | E | 201 | - | - | 12/41/41/41 | - |
| 19 | SF4 | C | 301 | 3 | - | - | 0/6/5/5 |
| 28 | FES | M | 803 | 12 | - | - | 0/1/1/1 |
| 28 | FES | O | 301 | 14 | - | - | 0/1/1/1 |

The worst 5 of 84 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|--------|-------------|----------|
| 21 | A | 503 | NAI | O4B-C1B | 16.12 | 1.63 | 1.41 |
| 21 | A | 503 | NAI | C2B-C1B | -15.32 | 1.30 | 1.53 |
| 26 | J | 401 | NDP | C3B-C2B | -12.78 | 1.24 | 1.52 |
| 26 | J | 401 | NDP | C6N-C5N | 12.40 | 1.55 | 1.33 |
| 26 | J | 401 | NDP | O4D-C4D | 10.75 | 1.69 | 1.45 |

The worst 5 of 60 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 26 | J | 401 | NDP | C3N-C2N-N1N | -7.99 | 111.69 | 123.10 |
| 27 | J | 402 | UQ | C7-C8-C9 | -7.64 | 114.07 | 126.79 |
| 26 | J | 401 | NDP | C1D-N1N-C2N | -6.37 | 110.50 | 121.11 |
| 24 | E | 201 | 8Q1 | C6-C1-S44 | 6.35 | 120.85 | 113.46 |
| 27 | J | 402 | UQ | C12-C13-C14 | -6.18 | 112.78 | 127.66 |

There are no chirality outliers.

5 of 136 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 20 | A | 502 | FMN | N10-C1'-C2'-O2' |
| 20 | A | 502 | FMN | N10-C1'-C2'-C3' |
| 20 | A | 502 | FMN | C3'-C4'-C5'-O5' |
| 20 | A | 502 | FMN | O4'-C4'-C5'-O5' |
| 20 | A | 502 | FMN | C5'-O5'-P-O1P |



There are no ring outliers.



12 monomers are involved in 29 short contacts:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 30 | Q | 501 | UQ1 | 1 | 0 |
| 21 | A | 503 | NAI | 5 | 0 |
| 25 | I | 201 | CDL | 1 | 0 |
| 23 | C | 303 | PLX | 3 | 0 |
| 30 | Q | 502 | UQ1 | 1 | 0 |
| 20 | A | 502 | FMN | 4 | 0 |
| 26 | J | 401 | NDP | 2 | 0 |
| 19 | A | 501 | SF4 | 2 | 0 |
| 24 | E | 201 | 8Q1 | 4 | 0 |
| 27 | J | 402 | UQ | 2 | 0 |
| 19 | C | 301 | SF4 | 1 | 0 |
| 22 | C | 302 | PEE | 4 | 0 |



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.


The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32210. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 222



Y Index: 210




Z Index: 200


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0162. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 124 nm^3 ; this corresponds to an approximate mass of 112 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.417\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-32210 and PDB model 7VYN. Per-residue inclusion information can be found in section 3 on page 15.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0162 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0162).

9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0162) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.8760 |  0.6810 |
| A |  0.8880 |  0.6720 |
| B |  0.9520 |  0.7140 |
| C |  0.9370 |  0.7090 |
| E |  0.8390 |  0.6730 |
| F |  0.6930 |  0.6130 |
| G |  0.3800 |  0.4880 |
| H |  0.8110 |  0.6490 |
| I |  0.7830 |  0.6440 |
| J |  0.8960 |  0.6870 |
| K |  0.7910 |  0.6380 |
| L |  0.8870 |  0.6940 |
| M |  0.9030 |  0.6910 |
| N |  0.8660 |  0.6840 |
| O |  0.8240 |  0.6510 |
| P |  0.9550 |  0.7130 |
| Q |  0.9660 |  0.7170 |
| T |  0.8530 |  0.6800 |
| W |  0.7410 |  0.6330 |

