



# Full wwPDB X-ray Structure Validation Report i

May 21, 2020 – 01:01 pm BST

PDB ID : 2Y7O  
Title : Structure of N-terminal domain of Candida albicans als9-2 - G299W mutant  
Authors : Salgado, P.S.; Burchell, L.; Cota, E.  
Deposited on : 2011-01-31  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	312	88%  8% ..

## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AGGLUTININ-LIKE ALS9 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	2362	1497	365	490	10	1	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP Q5A8T1
A	51	THR	ASN	conflict	UNP Q5A8T1
A	212	VAL	ILE	conflict	UNP Q5A8T1
A	296	TRP	GLY	engineered mutation	UNP Q5A8T1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
			Total	O	
2	A	275	275	275	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AGGLUTININ-LIKE ALS9 PROTEIN



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.37 Å   68.31 Å   89.35 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	36.51 – 1.75 46.46 – 1.26	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.51-1.75) 30.7 (46.46-1.26)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	7.39 (at 1.26 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
$R$ , $R_{free}$	0.195 , 0.223 0.550 , 0.522	Depositor DCC
$R_{free}$ test set	1639 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	4.4	Xtriage
Anisotropy	2.187	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.24 , 215.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.42	EDS
Total number of atoms	2637	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.74	14/2420 (0.6%)	1.19	4/3314 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	106	ARG	CZ-NH1	53.99	2.03	1.33
1	A	106	ARG	CZ-NH2	11.09	1.47	1.33
1	A	106	ARG	NE-CZ	9.77	1.45	1.33
1	A	301	GLU	CD-OE2	8.36	1.34	1.25
1	A	86	GLU	CD-OE1	6.43	1.32	1.25
1	A	234	TYR	CE1-CZ	-6.27	1.30	1.38
1	A	304	SER	C-O	6.18	1.35	1.23
1	A	24	LYS	CE-NZ	6.07	1.64	1.49
1	A	301	GLU	CD-OE1	5.94	1.32	1.25
1	A	265	GLU	CD-OE1	5.88	1.32	1.25
1	A	134	PHE	CE1-CZ	5.82	1.48	1.37
1	A	304	SER	CB-OG	5.76	1.49	1.42
1	A	143	PHE	CE2-CZ	5.58	1.48	1.37
1	A	188	CYS	CB-SG	-5.37	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	ARG	NE-CZ-NH2	-27.27	106.67	120.30
1	A	265	GLU	C-N-CA	-7.58	102.76	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	A	308	VAL	CG1-CB-CG2	-5.14	102.68	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	ARG	Sidechain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2362	0	2231	43	0
2	A	275	0	0	1	0
All	All	2637	0	2231	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:LYS:HA	1:A:299:ASN:CB	1.47	1.31
1:A:299:ASN:N	1:A:301:GLU:OE1	1.79	1.16
1:A:298:LYS:HA	1:A:299:ASN:HB2	1.18	1.14
1:A:301:GLU:O	1:A:302:ALA:HB3	1.39	1.10
1:A:298:LYS:HA	1:A:299:ASN:HB3	1.30	1.08
1:A:299:ASN:H	1:A:301:GLU:CD	1.61	1.02
1:A:298:LYS:CA	1:A:299:ASN:CB	2.38	1.02
1:A:303:ASP:CG	1:A:304:SER:H	1.66	0.97
1:A:298:LYS:NZ	1:A:300:SER:HB2	1.81	0.96
1:A:301:GLU:O	1:A:302:ALA:CB	2.12	0.93
1:A:298:LYS:CA	1:A:299:ASN:HB2	2.00	0.90
1:A:298:LYS:HG2	1:A:300:SER:N	1.87	0.90

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ASN:ND2	1:A:299:ASN:O	2.07	0.88
1:A:298:LYS:HZ3	1:A:300:SER:HB2	1.39	0.83
1:A:298:LYS:HZ3	1:A:300:SER:CB	1.94	0.81
1:A:17:ARG:HB3	2:A:2019:HOH:O	1.80	0.80
1:A:298:LYS:HG2	1:A:300:SER:H	1.45	0.79
1:A:303:ASP:CG	1:A:304:SER:N	2.40	0.75
1:A:300:SER:H	1:A:301:GLU:HA	1.52	0.74
1:A:298:LYS:NZ	1:A:300:SER:CB	2.50	0.74
1:A:298:LYS:C	1:A:301:GLU:OE1	2.28	0.71
1:A:298:LYS:CA	1:A:299:ASN:HB3	2.14	0.71
1:A:298:LYS:HD2	1:A:309:ILE:HD11	1.76	0.66
1:A:300:SER:N	1:A:301:GLU:HA	2.17	0.59
1:A:299:ASN:C	1:A:299:ASN:HD22	1.98	0.56
1:A:299:ASN:N	1:A:301:GLU:CD	2.44	0.55
1:A:298:LYS:O	1:A:301:GLU:OE1	2.25	0.55
1:A:303:ASP:OD2	1:A:304:SER:N	2.28	0.53
1:A:204:GLY:HA3	1:A:267:ILE:HD13	1.89	0.53
1:A:200:VAL:HG13	1:A:273:THR:HB	1.91	0.52
1:A:136:ALA:HB2	1:A:158:ALA:HB2	1.96	0.48
1:A:248:GLN:O	1:A:249:ASN:O	2.32	0.47
1:A:216:ILE:HG22	1:A:256:PRO:HA	1.99	0.44
1:A:24:LYS:HE3	1:A:24:LYS:HB3	1.85	0.44
1:A:298:LYS:HZ1	1:A:303:ASP:CB	2.31	0.44
1:A:136:ALA:CB	1:A:158:ALA:HB2	2.47	0.43
1:A:298:LYS:HZ1	1:A:303:ASP:HB3	1.83	0.43
1:A:298:LYS:CG	1:A:299:ASN:HB3	2.48	0.43
1:A:267:ILE:HD12	1:A:269:LYS:O	2.19	0.43
1:A:298:LYS:HZ2	1:A:300:SER:HB2	1.74	0.42
1:A:29:PRO:HA	1:A:30:THR:HA	1.88	0.41
1:A:298:LYS:NZ	1:A:303:ASP:HB2	2.36	0.41
1:A:248:GLN:O	1:A:249:ASN:C	2.57	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	309/312 (99%)	295 (96%)	10 (3%)	4 (1%)	12 2

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	249	ASN
1	A	299	ASN
1	A	305	ASP
1	A	303	ASP

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	272/274 (99%)	264 (97%)	8 (3%)	42 19

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	PHE
1	A	17	ARG
1	A	20	GLU
1	A	102	VAL
1	A	200	VAL
1	A	299	ASN
1	A	304	SER
1	A	305	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.