Full wwPDB NMR Structure Validation Report (i)

May 7, 2024 – 03:49 pm BST

PDB ID	:	3ZJ2
BMRB ID	:	18956
Title	:	Structure of Nab2p tandem zinc finger 34
Authors	:	Martinez-Lumbreras, S.; Santiveri, C.M.; Mirassou, Y.; Zorrilla, S.; Perez-Canadillas, J.M.
Deposited on	:	2013-01-16

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 95%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	71	76% • 23%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:338-A:392 (55)	0.41	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 6, 7, 8, 10, 11, 12, 15, 19, 20
2	4, 5, 13, 16, 17, 18
Single-model clusters	9; 14

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1069 atoms, of which 533 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	71	1067	332	533	93	102	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	331	GLY	-	expression tag	UNP P32505
A	332	SER	-	expression tag	UNP P32505

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
			Total	Zn
2	A	2	2	2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 76% · 23%

G331 S332 V333 Q334 T335 G336 I337 R383 K393 E394 V395 K396 P397 I398 S399 Q400 K401

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% · 23%

G331 S332 V333 Q334 T335 G336 I337 L375 K391 I392 K393 E394 V395 K396 P397 I398 S399 Q400 K401

4.2.2 Score per residue for model 2

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% · 23%

G331 S332 V333 Q334 T335 G336 I337 S388 L389 S390 K393 E394 V395 K396 P397 I398 S399 Q400 K401

4.2.3 Score per residue for model 3

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 72% 23%

4.2.4 Score per residue for model 4

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% 23%



4.2.5 Score per residue for model 5

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 76% 23%



4.2.6 Score per residue for model 6

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 77% 23%



4.2.7 Score per residue for model 7

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 70% 6% 23%

4.2.8 Score per residue for model 8 (medoid)

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A:

4.2.9 Score per residue for model 9

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A:

4.2.10 Score per residue for model 10

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A:



4.2.11 Score per residue for model 11

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A:

4.2.12 Score per residue for model 12

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A:



4.2.13 Score per residue for model 13

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% • 23%



4.2.14 Score per residue for model 14

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 72% 6% 23%

4.2.15 Score per residue for model 15

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 72% 6% 23%

4.2.16 Score per residue for model 16

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 76% • 23%



4.2.17 Score per residue for model 17

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% • 23%



4.2.18 Score per residue for model 18

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 76% • 23%



4.2.19 Score per residue for model 19

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 75% • 23%



4.2.20 Score per residue for model 20

• Molecule 1: NUCLEAR POLYADENYLATED RNA-BINDING PROTEIN NAB2

Chain A: 72% 6% 23%

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *CYANA, AMBER*.

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *LOWEST TARGET FUNCTION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	7.0
NMRPipe	structure solution	
CcpNmr Analysis	structure solution	
CYANA	structure solution	
Amber	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	859
Number of shifts mapped to atoms	859
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	95%

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.62±0.01	0±0/426 (0.0± 0.0%)	1.01±0.02	1±1/579 (0.1± 0.1%)
All	All	0.62	0/8520 (0.0%)	1.01	14/11580 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.2
All	All	0	1

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	383	ARG	NE-CZ-NH1	6.28	123.44	120.30	3	8
1	A	383	ARG	NE-CZ-NH2	-5.84	117.38	120.30	19	6

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	383	ARG	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	415	400	400	0±0
All	All	8340	8000	8000	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:392:ILE:HD12	1:A:392:ILE:H	0.46	1.71	11	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	55/71 (77%)	49±1 (89±2%)	6±1 (11±2%)	0±0 (0±0%)	54	85
All	All	1100/1420 (77%)	982 (89%)	117 (11%)	1 (0%)	54	85

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	341	LYS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	49/63 (78%)	47±1 (96±3%)	2±1 (4±3%)	32 81
All	All	980/1260 (78%)	937 (96%)	43 (4%)	32 81

All 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	375	LEU	10
1	A	383	ARG	5
1	A	361	GLU	4
1	A	389	LEU	4
1	A	391	LYS	3
1	A	388	SER	3
1	A	373	LYS	3
1	A	390	SER	2
1	A	341	LYS	2
1	A	369	MET	2
1	A	392	ILE	2
1	A	368	LEU	1
1	A	364	LYS	1
1	A	339	LEU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 95% for the well-defined parts and 94% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *Nab2_Z34_CS_curated.dat.csh*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	859
Number of shifts mapped to atoms	859
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	71	0.03 ± 0.20	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	67	0.39 ± 0.16	None needed (< 0.5 ppm)
$^{13}C'$	69	0.22 ± 0.19	None needed (< 0.5 ppm)
^{15}N	63	-1.23 ± 0.52	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 95%, i.e. 664 atoms were assigned a chemical shift out of a possible 698. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	267/267 (100%)	107/107 (100%)	110/110 (100%)	50/50 (100%)
Sidechain	359/383 (94%)	244/249 (98%)	111/122 (91%)	4/12 (33%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	38/48 (79%)	22/24 (92%)	11/19 (58%)	5/5 (100%)
Overall	664/698 (95%)	373/380 (98%)	232/251 (92%)	59/67 (88%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 94%, i.e. 856 atoms were assigned a chemical shift out of a possible 911. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	339/347 (98%)	136/140 (97%)	140/142 (99%)	63/65 (97%)
Sidechain	479/516 (93%)	325/335 (97%)	150/164 (91%)	4/17 (24%)
Aromatic	38/48 (79%)	22/24 (92%)	11/19 (58%)	5/5 (100%)
Overall	856/911 (94%)	483/499 (97%)	301/325 (93%)	72/87 (83%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	357	THR	HG1	5.61	0.08 – 2.19	21.2
1	A	370	TRP	CZ2	132.03	107.20 – 121.33	12.6

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:


RCI
0.0
0.1
0.2
0.3
0.4
0.5
331:GLY
334:GLN
337:ILE
340:CYS
343:GLY
346:CYS
349:PRO
352:PRO
355:HIS
358:PRO
361:GLU
364:LYS
367:ASP
370:TRP
373:LYS
376:THR
379:ASN
382:CYS
385:ALA
388:SER
391:LYS
394:GLU
397:PRO
400:GLN
Auth_seq_ID